

Introduction

A worker honey bee performs different tasks throughout its life span. The shift from one task to another, especially from in-hive tasks to outside tasks requires partly drastic adaptations to a different environment. One major difference between the in-hive environment and the outside is exposure to light as bees leave the dark hive (Fig. 1).

Light plays an essential role in the life of the forager bees in terms of visual navigation and spotting of food sources. It is apparent that for these important tasks foragers need to be optimally prepared by adaptive changes in the neuronal circuitry. Indeed, the transition from in-hive tasks to foraging is associated with remarkable changes in brain structure, and with synaptic plasticity¹. Even the exposure of adult worker bees to artificial light is sufficient to induce structural synaptic plasticity in visual subcompartments of the mushroom bodies².

To investigate how synaptic plasticity in visual centers of the brain is controlled at the level of genes, we compared gene expression levels in the optic lobes of light exposed and dark kept honey bees via whole transcriptome sequencing (RNAseq) and quantitative real time PCR (qPCR).




Fig 1: Life of a worker honey bee

Methods

Procedure

- Bees were collected directly after eclosion and were kept in cages in total darkness.
- At the age of 1 or 7 days the bees were exposed to 5 x 45 min pulses of artificial full-spectrum day-light whereas an age-matched control group was kept in darkness (Fig. 2).
- The bees were snap frozen directly after the last pulse of the light program (Fig. 2).
- Optic lobes of 5 or 3 brains were dissected and pooled for RNAseq (n=2) and qPCR (n=8) analysis (Fig. 3).

Light program, sampling point




Fig 2: Light regime and sampling points

Brain dissection and pooling




Fig 3: Brain dissection and pooling

RNAseq with 1 day old bees

→ Identification of differentially expressed genes in the optic lobes. See top 52 in Tab 1.

Validation with qPCR

→ Of so far 7 tested genes 3 could be confirmed for 1 and 7 day old bees by qPCR: *CNPY-1-like* (GB50831), *Ip3k1* (GB41220), *Trim71* (GB48462).

→ *Trim71* is regulated by the micro RNA let-7³.



Results

| Gene (BeeBase) | Light induced expression | Reproducibility in experiment 1 and 2 | General function |
|----------------|--------------------------|---------------------------------------|----------------------------------|
| GB55613* | Up | YES | Unknown |
| GB54595* | Down | YES | histone H3-K27 demethylase |
| GB45148 | Up | YES | Vitamin A related |
| GB45147* | UP | YES | Vitamin A related |
| GB45024 | UP | YES | Vitamin A related |
| GB45023 | Up | YES | Vitamin A related |
| GB41220* | Up | YES | IP3 kinase |
| GB42985 | Up | Weaker induction in 2 | pyruvate lyase |
| GB41002 | Up | YES | timeless |
| GB43805 | Up | YES | metallo-endopeptidase |
| GB46312 | Up | Weaker induction in 2 | cuticular protein |
| GB55396 | Up | YES | Unknown |
| GB50831* | Up | YES | neurite outgrowth enhancer |
| GB48462 | Up | YES | E3 ubiquitin protein ligase |
| GB43732 | Up | YES | serine/threonine-protein kinase |
| GB44871 | Up | YES | glycine N-methyltransferase |
| GB47279 | Up | YES | cytochrome P450 |
| GB43514 | Up | YES | lipase, member H |
| GB49843 | Up | YES | neuronal PAS domain protein |
| GB54962 | Up | YES | Unknown |
| GB42197 | Up | YES | Unknown |
| GB47484 | Up | YES | Histone-like? |
| GB47382 | Up | YES | Histone H4 |
| GB41720 | Up | YES | Pleckstrin-like |
| GB48492* | UP | YES | Take-out |
| GB42467 | UP | YES | phototransduction |
| GB42673 | UP | Weaker induction in 1 | RDH10/retinol dehydrogenase |
| GB43649 | UP | YES | chloride channel |
| GB55043 | UP | YES | kainate glutamate receptor |
| GB43823* | UP | YES | chemosensory protein CSP1 |
| GB41593* | UP | YES | cell migration regulator |
| GB40046 | UP | YES | neuronal mt transport protein |
| GB55050 | UP | YES | transmembrane transporter |
| GB41277* | UP | YES | light-induced ubiquitylation |
| GB45365 | UP | YES | transmembrane transporter |
| GB47948 | UP | YES | myosin light chain kinase |
| GB41720 | UP | YES | Pleckstrin-like |
| GB51220 | UP | YES | cytochrome b-561 |
| GB40552 | UP | YES | Unknown |
| GB45910 | UP | YES | Crystallin |
| GB45906 | UP | YES | Crystallin 2 |
| GB46514 | UP | YES | Acetylcholinesterase (both loci) |
| GB44095 | UP | YES | Cation channel |
| GB42227 | UP | Weaker induction in 1 | Homeobox related |
| GB51580* | UP | YES | acyl-CoA synthetase |
| GB41339 | UP | Weaker induction in 2 | acid phosphatase |
| GB52448 | UP | YES | Unknown |
| GB53210 | UP | YES | Unknown |
| GB47697 | UP | YES | Unknown (both loci) |
| GB47697 | UP | YES | Unknown (both loci) |
| GB41709 | UP | YES | Unknown |

Table 1: Top 52 differentially expressed genes in the optic lobes identified by RNAseq
These 52 genes show the biggest difference in their expression level between 1 day old light exposed-, and 1 day old dark kept bees. Genes with an epigenetic function are highlighted in green; *methylated genes;

Fig 4: Relative gene expression ratios (R) by qPCR analysis

Relative gene expression ratios (R) were calibrated against 1 day old dark kept bees (1d Dark). (A) shows R for *Trim71* (GB48462), and (B) shows R for its negative regulator micro RNA let-7. 1d: 1 day old bees; 7d: 7 day old bees;



Conclusion and Outlook

- RNAseq revealed genes which show a difference in their expression levels in the honey bees' optic lobes between the two treatment groups (Tab. 1).
- Several of these genes (i.e. *CNPY-1-like* (GB50831), *Ip3k1* (GB41220)) have been associated with neuronal plasticity and therefore are good candidates to play a role in light induced synaptic plasticity in the honey bee brain^{4,5}.
- Some genes like *Trim71* (GB48462) and *histone demethylase UTY-like* (GB54595) could rather contribute to an adaptation to a changed environment by epigenetic mechanisms which influence the transcription of a broad number of genes simultaneously.
- qPCR studies could confirm some of the tested genes found with RNAseq, but failed to confirm others. This might be due to the partly different bee handling and different honey bee races used in both applications as RNAseq was performed in Canberra, Australia with *A.m. var. ligustica*, whereas qPCR experiments took place in Würzburg, Germany with *A.m. var. carnica*.
- Future studies will aim at finding a causality between expression of the candidate genes, synaptic plasticity and subsequent behavior. For this issue pharmacological or siRNA knockdown experiments are planned.