

References

Abe, K. and Sugita, Y. (1979) Properties of cytochrome *b*₅ and methemoglobin reduction in human erythrocytes. *Eur. J. Biochem.* **101**, 423-428.

Adachi, K. (1972) Studies on reduced pyridine nucleotide dehydrogenase in bovine erythrocytes. II. Electron acceptor specificity of two types of reduced pyridine nucleotide dehydrogenase in bovine erythrocytes. *Biochim. Biophys. Acta* **289**, 262-268.

Agar, N.S., Reinke, N.B., Godwin, I.R. and Kuchel, P.W. (2000) Comparative biochemistry of marsupial erythrocytes: A review. *Comp. Haematol. Int.* **10**, 148-167.

Agutter, P.S., Sherriff, R.V. and Kadlubowski, M. (1979) A novel lipid hydroperoxide-reducing enzyme in human erythrocyte membranes. *Biochem. Soc. Trans.* **7**, 710-711.

Agutter, P.S., Dallas, J.M., Newman, P. and Kadlubowski, M. (1980) Evidence for the existence of two NADH-ferricyanide reductases in human erythrocyte membranes. *Biochem. Soc. Trans.* **8**, 326.

Aoshiha, K., Nakajima, Y., Yasui, S., Tamaoki, J. and Nagai, A. (1999) Red blood cells inhibit apoptosis of human neutrophils. *Blood* **93**, 4006-4010.

Augustyniak, K., Zavodnik, I.B., Palecz, D., Szosland, K. and Bryszewska, M. (1996) The effect of oxidising agents and diabetes mellitus on the human red blood cell membrane potential. *Clin. Biochem.* **29**, 283-286.

- Avron, M. and Shavit, N. (1963) A sensitive and simple determination of ferrocyanide. *Anal. Biochem.* **6**, 549-554.
- Baker, M.A. and Lawen, A. (2000) The function of the plasma membrane NADH-oxidoreductase system. A critical review of the structural and functional data. *Antioxid. Redox. Signaling* **2**, 197-212.
- Baker, M.A., Krutskikh, A., Curry, B.J., McLaughlin, E.A. and Aitken, R.J. (2004a) Identification of cytochrome *P*450-reductase as the enzyme responsible for NADPH-dependent lucigenin and tetrazolium salt reduction in rat epididymal sperm preparations. *Biol. Reprod.* **71**, 307-318.
- Baker, M.A., Lane, D.J.R., Ly, J.D., De Pinto, V. and Lawen, A. (2004b) VDAC1 is a transplasma membrane NADH-ferricyanide reductase. *J. Biol. Chem.* **279**, 4811-4819.
- Berridge, M.V. and Tan, A.S. (1993) Characterization of the cellular reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT): subcellular localization, substrate dependence, and involvement of mitochondrial electron transport in MTT reduction. *Arch. Biochem. Biophys.* **303**, 474-482.
- Berridge, M.V., Tan, A.S., McCoy, K.D. and Wang, R. (1996) The biochemical and cellular basis of cell proliferation assays that use tetrazolium salts. *Biochemica* **4**, 15-20.
- Berridge, M.V. and Tan, A.S. (1998) Trans-plasma membrane electron transport: a cellular assay for NADH- and NADPH-oxidase based on extracellular, superoxide-mediated reduction of the sulfonated tetrazolium salt WST-1. *Protoplasma* **205**, 74-82.
- Berridge, M.V. and Tan, A.S. (2000a) High capacity redox control at the plasma membrane of mammalian cells: trans membrane, cell-surface and serum NADH-oxidases. *Antioxid. Redox. Signaling* **2**, 231-242.

- Berridge, M.V. and Tan, A.S. (2000b) Cell surface NAD(P)H-oxidase: relationship to transplasma membrane NADH-oxidoreductase and a potential source of circulating NADH-oxidase. *Antioxid. Redox. Signaling* **2**, 277-288.
- Berthon, H.A. and Kuchel, P.W. (1995) NMR studies of erythrocyte metabolism. In *Advances in Molecular and Cell Biology, VII* (Brindle, K.M., Ed.). pp 147-205, JAI Press Inc., London.
- Beutler, E. (1984) Red Cell Metabolism - a Manual of Biochemical Methods. Grune & Stratton, Orlando.
- Birkmayer, J.G. (1996) Coenzyme nicotinamide adenine dinucleotide: new therapeutic approach for improving dementia of the Alzheimer type. *Ann. Clin. Lab. Sci.* **26**, 1-9.
- Birkmayer, J.G.D., Vrecko, C., Volc, D. and Birkmayer, W. (1993) Nicotinamide adenine-dinucleotide (NADH) - a new therapeutic approach to Parkinson's-disease - comparison of oral and parenteral application. *Acta Neurol. Scand.* **87**, 32-35.
- Bozzi, A., Parisi, M. and Strom, R. (1996) Erythrocyte glutathione determination in the diagnosis of glucose-6-phosphate dehydrogenase deficiency. *Biochem. Mol. Biol. Int.* **40**, 561-569.
- Brightman, A.O., Wang, J., Miu, R.K., Sun, I.L., Barr, R., Crane, F.L. and Morr , D.J. (1992) A growth factor- and hormone-stimulated NADH oxidase from rat liver plasma membrane. *Biochim. Biophys. Acta* **1105**, 109-117.
- Brindle, K.M., Brown, F.F., Campbell, I.D., Grathwohl, C. and Kuchel, P.W. (1979) Application of spin-echo nuclear magnetic resonance to whole-cell systems. Membrane transport. *Biochem. J.* **180**, 37-44.

- Brindle, K.M., Campbell, I.D. and Simpson, R.J. (1982) A ^1H NMR study of the kinetic properties expressed by glyceraldehyde phosphate dehydrogenase in the intact human erythrocyte. *Biochem. J.* **208**, 583-592.
- Brown, F.F., Campbell, I.D., Kuchel, P.W. and Rabenstein, D.C. (1977) Human erythrocyte metabolism studies by ^1H spin echo NMR. *FEBS Lett.* **82**, 12-16.
- Bruder, G., Bretscher, A., Franke, W.W. and Jarasch, E.D. (1980) Plasma membranes from intestinal microvilli and erythrocytes contain cytochromes b_5 and P-420. *Biochim. Biophys. Acta* **600**, 739-755.
- Bulliard, C., Zurbriggen, R., Tornare, J., Faty, M., Dastoor, Z. and Dreyer, J.L. (1997) Purification of a dichlorophenol-indophenol oxidoreductase from rat and bovine synaptic membranes: tight complex association of a glyceraldehyde-3-phosphate dehydrogenase isoform, TOAD64, enolase- γ and aldolase C. *Biochem. J.* **324**, 555-563.
- Chang, S.H. and Low, P.S. (2001) Regulation of the glycophorin C-protein 4.1 membrane-to-skeleton bridge and evaluation of its contribution to erythrocyte membrane stability. *J. Biol. Chem.* **276**, 22223-22230.
- Chaudiere, J. and Ferrari-Iliou, R. (1999) Intracellular antioxidants: from chemical to biochemical mechanisms. *Food Chem. Toxicol.* **37**, 949-962.
- Choury, D., Wajcman, H., Boissel, J.P. and Kaplan, J.C. (1981) Evidence for endogenous proteolytic solubilization of human red-cell membrane NADH-cytochrome b_5 reductase. *FEBS Lett.* **126**, 172-174.
- Chueh, P.J. (2000) Cell membrane redox systems and transformation. *Antioxid. Redox. Signaling* **2**, 177-187.
- Chueh, P.J., Kim, C., Cho, N., Morr , D.M. and Morr , D.J. (2002) Molecular cloning and characterization of a tumor-associated, growth-related, and time-keeping

- hydroquinone (NADH) oxidase (tNOX) of the HeLa cell surface. *Biochemistry* **41**, 3732-3741.
- Ciriolo, A.R., Paci, M., Sette, M., de Martino, A., Bozzi, A. and Rotilio, G. (1993) Transduction of reducing power across the plasma membrane by reduced glutathione. *Eur. J. Biochem.* **215**, 711-718.
- Ciriolo, M.R., Sette, M., Paci, M. and Rotilio, G. (1990) A study of the intracellular effects of glutathione by H-1-spin echo NMR of intact human erythrocytes. *Biochem. Int.* **20**, 397-403.
- Clark, M.G., Partick, E.J. and Crane, F.L. (1982) Properties and regulation of a trans-plasma membrane redox system in rat liver. *Biochem. J.* **204**, 795-801.
- Cochet, S., Volet, G., Cartron, J.P. and Bertrand, O. (2001) New procedures for glycophorin A purification with high yield and high purity. *J. Chromatogr. B. Biomed. Sci. App.* **750**, 109-119.
- Constantinescu, A., Han, D. and Packer, L. (1993) Vitamin E recycling in human erythrocyte membranes. *J. Biol. Chem.* **268**, 10906-10913.
- Cotgreave, I.A. and Gerdes, R.G. (1998) Recent trends in glutathione biochemistry--glutathione-protein interactions: a molecular link between oxidative stress and cell proliferation? *Biochem. Biophys. Res. Commun.* **242**, 1-9.
- Crane, F.L., Crane, H.E., Sun, I.L., MacKellar, W.C., Grebing, C. and Löw, H. (1982) Insulin control of a transplasma membrane NADH dehydrogenase in erythrocyte membranes. *J. Bioenerg. Biomembr.* **14**, 425-433.
- Crane, F.L., Sun, I.L., Crowe, R.A., Alcain, F.J. and Löw, H. (1994) Coenzyme Q(10), plasma-membrane oxidase and growth-control. *Mol. Aspects Med.* **15**, 1-11.

Dancey, G.F. and Shapiro, B.M. (1976) The NADH dehydrogenase of the respiratory chain of *Escherichia coli*. II. Kinetics of the purified enzyme and the effects of antibodies elicited against it on membrane-bound and free enzyme. *J. Biol. Chem.* **251**, 5921-5928.

Davies, K.J. and Goldberg, A.L. (1987) Oxygen radicals stimulate intracellular proteolysis and lipid peroxidation by independent mechanisms in erythrocytes. *J. Biol. Chem.* **262**, 8220-8226.

Dawson-Saunders, B. and Trapp, R.G. (1994) *Basic and Clinical Biostatistics*. Appleton & Lange, Connecticut.

Debnam, P.M. and Shearer, G. (1997) Colorimetric assay for substrates of NADP⁺-dependent dehydrogenases based on reduction of a tetrazolium dye to its soluble formazan. *Anal. Biochem.* **250**, 253-255.

del Castillo-Olivares, A., Marquez, J., Nunez De Castro, I. and Medina, M.A. (1994) Characterization of plasma membrane redox activity from Ehrlich cells. *Cell Biochem. Funct.* **12**, 149-152.

del Castillo-Olivares, A., Medina, M.A., Nunez de Castro, I. and Marquez, J. (1996) Purification and characterization of a plasma membrane ferricyanide-utilizing NADH dehydrogenase from Ehrlich tumor cells. *Biochem. J.* **314**, 587-593.

del Castillo-Olivares, A., de Castro, I.N. and Medina, M.A. (2000) Dual role of plasma membrane electron transport systems in defense. *Crit. Rev. Biochem. Mol. Biol.* **35**, 197-220.

Demehin, A.A., Abugo, O.O. and Rifkind, J.M. (2001) The reduction of nitroblue tetrazolium by red blood cells: a measure of red cell membrane antioxidant capacity and hemoglobin-membrane binding sites. *Free Radic. Res.* **34**, 605-620.

- Dierickx, P.J., Nuffel, G.V. and Alvarez, I. (1999) Glutathione protection against hydrogen peroxide, *tert*-butyl hydroperoxide and diamide cytotoxicity in rat hepatoma-derived Fa32 cells. *Hum. Exp. Toxicol.* **18**, 627-633.
- Dormandy, T.L. and Zarday, Z. (1965) The mechanism of insulin action: the immediate electrochemical effects of insulin on red-cell systems. *J. Physiol.* **180**, 684-707.
- Dotsch, J., Demirakca, S., Cryer, A., Hanze, J., Kuhl, P.G. and Rascher, W. (1998) Reduction of NO-induced methemoglobinemia requires extremely high doses of ascorbic acid in vitro. *Intensive Care Med.* **24**, 612-615.
- Dreyer, J.L. (1990) Plasma membrane dehydrogenase in rat brain synaptic membranes. Multiplicity and subunit composition. *J. Bioenerg. Biomembr.* **22**, 619-633.
- Ellem, K.A. and Kay, G.F. (1983) Ferricyanide can replace pyruvate to stimulate growth and attachment of serum restricted human melanoma cells. *Biochem. Biophys. Res. Commun.* **112**, 183-190.
- Falke, J.J. and Chan, S.I. (1986) Molecular mechanisms of band 3 inhibitors. 1. Transport site inhibitors. *Biochemistry* **25**, 7888-7894.
- Fehlau, R., Grygorczyk, R., Fuhrmann, G.F. and Schwarz, W. (1989) Modulation of the Ca²⁺- or Pb²⁺-activated K⁺-selective channels in human red cells. II. Parallelisms to modulation of the activity of a membrane-bound oxidoreductase. *Biochim. Biophys. Acta* **978**, 37-42.
- Ferreira, A.L., Machado, P.E. and Matsubara, L.S. (1999) Lipid peroxidation, antioxidant enzymes and glutathione levels in human erythrocytes exposed to colloidal iron hydroxide *in vitro*. *Brazil. J. Med. Biol. Res.* **32**, 689-694.

- Filomeni, G., Rotilio, G. and Ciriolo, M. (2002) Cell signaling and the glutathione redox system. *Biochem. Pharmacol.* **64**, 1057-1064.
- Fiorani, M., De Sanctis, R., De Bellis, R. and Dachà, M. (2002) Intracellular flavonoids as electron donors for extracellular ferricyanide reduction in human erythrocytes. *Free Radic. Biol. Med.* **32**, 64-72.
- Foldes-Papp, Z. and Mareyzki, D. (1984) Enzymatic *t*-butyl hydroperoxide reduction on human erythrocyte membranes - NADPH and GSH dependent activities. *Biomed. Biochim. Acta* **3**, 271-279.
- Forsyth, L.M., Preuss, H.G., MacDowell, A.L., Chiazze, L., Birkmayer, G.D. and Bellanti, J.A. (1999) Therapeutic effects of oral NADH on the symptoms of patients with chronic fatigue syndrome. *Ann. Allergy. Asthma. Immunol.* **82**, 185-191.
- Fridovich, I. (1997) Superoxide anion radical ($O_2^{\cdot-}$), superoxide dismutases, and related matters. *J. Biol. Chem.* **272**, 18515-18517.
- Giardina, B., Messina, I., Scatena, R. and Castagnola, M. (1995) The multiple functions of hemoglobin. *Crit. Rev. Biochem. Mol. Biol.* **30**, 165-196.
- Goldenberg, H., Crane, F.L. and Morrè, D.J. (1978) Influence of hormones on NADH-dehydrogenase in mouse liver plasma membrane. *Biochem. Biophys. Res. Commun.* **83**, 234-240.
- Goldenberg, H., Crane, F.L. and Morrè, D.J. (1979) NADH oxidoreductase of mouse liver plasma membranes. *J. Biol. Chem.* **254**, 2491-2498.
- Goldenberg, H., Landertshamer, H. and Laggner, H. (2000) Functions of vitamin C as a mediator of transmembrane electron transport in blood cells and related cell culture models. *Antioxid. Redox. Signaling* **2**, 189-196.

- Gomez-Diaz, C., Rodriguez-Aguilera, J.C., Barroso, M.P., Villalba, J.M., Navarro, F., Crane, F.L. and Navas, P. (1997) Antioxidant ascorbate is stabilized by NADH-coenzyme Q10 reductase in the plasma membrane. *J. Bioenerg. Biomembr.* **29**, 251-257.
- Grebing, C., Crane, F.L., Löw, H. and Hall, K. (1984) A transmembranous NADH-dehydrogenase in human erythrocyte membranes. *J. Bioenerg. Biomembr.* **16**, 517-533.
- Greiber, S., Münzel, T., Kästner, S., Müller, B., Schollmeyer, P. and Pavenstädt, H. (1998) NAD(P)H oxidase activity in cultured human podocytes: effects of adenosine triphosphate. *Kidney Int.* **53**, 654-663.
- Guarnieri, C., Fraticelli, A., Ventura, C., Vaona, I. and Budini, R. (1987) External GSSG enhances intracellular glutathione in isolated cardiac myocytes. *Biochem. Biophys. Res. Commun.* **147**, 658-665.
- Günther, H. (1980) *NMR Spectroscopy An Introduction*. John Wiley & Sons, New York.
- Halestrap, A.P. and Denton, R.M. (1974) Specific inhibition of pyruvate transport in rat liver mitochondria and human erythrocytes by alpha-cyano-4-hydroxycinnamate. *Biochem. J.* **138**, 313-316.
- Halestrap, A.P. and Denton, R.M. (1975) The specificity and metabolic implications of the inhibition of pyruvate transport in isolated mitochondria and intact tissue preparations by alpha-cyano-4-hydroxycinnamate and related compounds. *Biochem. J.* **148**, 97-106.
- Halestrap, A.P. (1976) Transport of pyruvate and lactate into human erythrocytes. Evidence for the involvement of the chloride carrier and a chloride-independent carrier. *Biochem. J.* **156**, 193-207.

Harkness, D.R., Ponce, J. and Grayson, V. (1969) A comparative study on the phosphoglyceric acid cycle in mammalian erythrocytes. *Comp. Biochem. Physiol.* **28**, 129-138.

Harvey, J.W. (1989) Erythrocyte metabolism. In *Clinical Biochemistry of Domestic Animals, 4th edn* (Kaneko, J.J., Ed.). pp 185-233, Academic Press, New York.

Hedges, K.L., Morr , D.M., Wu, L.Y. and Morr , D.J. (2003) Adriamycin tolerance in human mesothelioma lines and cell surface NADH oxidase. *Life Sci.* **73**, 1189-1198.

Hegesh, E., Gruener, N., Cohen, S., Bochkovsky, R. and Shuval, H.I. (1970) A sensitive micromethod for the determination of methemoglobin in blood. *Clin. Chim. Acta* **30**, 679-682.

Himmelreich, U. and Kuchel, P.W. (1997) ¹³C-NMR studies of transmembrane electron transfer to extracellular ferricyanide in human erythrocytes. *Eur. J. Biochem.* **246**, 638-645.

Himmelreich, U., Drew, K., Serianni, A.S. and Kuchel, P.W. (1998) ¹³C NMR studies of vitamin C transport and its redox cycling in human erythrocytes. *Biochemistry* **37**, 7578-7588.

Hsieh, H.S. and Jaff , E.R. (1975) The metabolism of methemoglobin in human erythrocytes. In *The Red Blood Cell* (Surgenor, D.M.N., Ed.). pp 799-824, Academic Press II, New York.

Hubbard, A.R., Sprandel, U. and Chalmers, R.A. (1980) Organic-acid transport in resealed haemoglobin-containing human erythrocyte 'ghosts'. *Biochem. J.* **190**, 653-658.

- Ingrosso, D., D'Angelo, S., di Carlo, E., Perna, A.F., Zappia, V. and Galletti, P. (2000) Increased methyl esterification of altered aspartyl residues in erythrocyte membrane proteins in response to oxidative stress. *Eur. J. Biochem.* **267**, 4397-4405.
- Inman, R.S. and Wessling-Resnick, M. (1993) Characterization of transferrin-independent iron transport in K562 cells. Unique properties provide evidence for multiple pathways of iron uptake. *J. Biol. Chem.* **268**, 8521-8528.
- Inman, R.S., Coughlan, M.M. and Wessling-Resnick, M. (1994) Extracellular ferrireductase activity of K562 cells is coupled to transferrin-independent iron transport. *Biochemistry* **33**, 11850-11857.
- Ishiyama, M., Shiga, M., Sasamoto, K., Mizoguchi, M. and He, P.G. (1993) A new sulfonated tetrazolium salt that produces a highly water-soluble formazan dye. *Chem. Pharm. Bull.* **41**, 1118-1122.
- Jaffé, E.R. (1981) Methaemoglobinaemia. *Clin. Haematol.* **10**, 99-122.
- Jaffé, E.R. and Hultquist, D.E. (1995) Cytochrome *b*₅ reductase deficiency and enzymopenic hereditary methemoglobinemia. In *The Metabolic and Molecular Bases of Inherited Disease* (Stanbury, J.B., Wyngaarden, J.B. and Fredrickson, D.G., Ed.). pp 2269-2280, McGraw-Hill, New York.
- Jansson, I. and Schenkman, J.B. (1977) Studies on three microsomal electron transfer enzyme systems. Specificity of electron flow pathways. *Arch. Biochem. Biophys.* **178**, 89-107.
- Jarvis, S.M., Young, J.D., Ansay, M., Archibald, A.L., Harkness, R.A. and Simmonds, R.J. (1980) Is inosine the physiological energy source of pig erythrocytes? *Biochim. Biophys. Acta* **597**, 183-188.
- Jeevanandam, M., Begay, C.K., Shahbazian, L.M. and Petersen, S.R. (2000) Altered plasma cytokines and total glutathione levels in parenterally fed critically ill trauma

patients with adjuvant recombinant human growth hormone (rhGH) therapy. *Crit. Care Med.* **28**, 324-329.

Jennings, M.L. (1976) Proton fluxes associated with erythrocyte membrane anion exchange. *J. Membr. Biol.* **28**, 187-205.

Jin, J.S., Webb, R.C. and D'Alecy, L.G. (1995) Inhibition of vascular nitric oxide-cGMP pathway by plasma from ischemic hindlimb of rats. *Am. J. Physiol.* **269**, H254-261.

Jozwik, M., Jozwik, M., Jozwik, M., Szczypka, M., Gajewska, J. and Laskowska-Klita, T. (1997) Antioxidant defence of red blood cells and plasma in stored human blood. *Clin. Chim. Acta* **267**, 129-142.

Kalinowski, H.O., Berger, S. and Braun, S. (1995) *Carbon-13 NMR Spectroscopy*. John Wiley & Sons, Chichester.

Kaneko, J.J., Tanaka, S., Nakajima, H. and Ushimi, C. (1969) Enzymes of equine erythrocytes: changes during equine infectious anemia. *Am. J. Vet. Res.* **30**, 543-549.

Kaneko, J.J. (1973) *Standard Values in Domestic Animals*, University of California, Davis

Kennett, E.C. and Kuchel, P.W. (2003) Redox reactions and electron transfer across the red cell membrane. *IUBMB Life* **55**, 375-385.

Kim, C., Crane, F.L., Becker, G.W. and Morr , D.J. (1995) Purification of NADH-cytochrome *b*₅ reductase from rat liver plasma membranes. *Protoplasma* **184**, 111-117.

Kim, C., Crane, F.L., Faulk, W.P. and Morr , D.J. (2002) Purification and characterization of a doxorubicin-inhibited NADH-quinone (NADH-ferricyanide) reductase from rat liver plasma membranes. *J. Biol. Chem.* **277**, 16441-16447.

- Kirk, K. and Kuchel, P.W. (1988a) Physical basis of the effect of hemoglobin on the ^{31}P NMR chemical shifts of various phosphoryl compounds. *Biochemistry* **27**, 8803-8810.
- Kirk, K. and Kuchel, P.W. (1988b) Characterization of transmembrane chemical shift differences in the ^{31}P NMR spectra of various phosphoryl compounds added to erythrocyte suspensions. *Biochemistry* **27**, 8795-8802.
- Kishi, T., Morr , D.M. and Morr , D.J. (1999) The plasma membrane NADH oxidase of HeLa cells has hydroquinone oxidase activity. *Biochim. Biophys. Acta* **1412**, 66-77.
- Kitajima, S., Yasukochi, Y. and Minakami, S. (1981) Purification and properties of human erythrocyte membrane NADH-cytochrome b_5 reductase. *Arch. Biochem. Biophys.* **210**, 330-339.
- Kuchel, P.W., Chapman, B.E., Endre, Z.H., King, G.F., Thorburn, D.R. and York, M.J. (1984) Monitoring metabolic reactions in erythrocytes using NMR spectroscopy. *Biomed. Biochim. Acta* **43**, 719-726.
- Kuchel, P.W. and Chapman, B.E. (1985) Direct observation of the NAD glycohydrolase reaction in human erythrocytes using NMR spectroscopy. *Experientia* **41**, 53-55.
- Kuchel, P.W. (1989) Biological applications of NMR. In *Analytical NMR* (Field, L.D. and Sternhell, S., Ed.). pp 157-219, John Wiley & Sons Ltd,
- Lapshina, E.A. and Bartosz, G. (1995) Nitrite-induced export of glutathione disulfide from erythrocytes. *Biochem. Mol. Biol. Int.* **37**, 949-957.

- Larm, J.A., Vaillant, F., Linnane, A.W. and Lawen, A. (1994) Up-regulation of the plasma membrane oxidoreductase as a prerequisite for the viability of human Namalwa ρ^0 cells. *J. Biol. Chem.* **269**, 30097-30100.
- Lawen, A., Martinus, R.D., McMullen, G.L., Nagley, P., Vaillant, F., Wolvetang, E.J. and Linnane, A.W. (1994) The universality of bioenergetic disease - the role of mitochondrial mutation and the putative interrelationship between mitochondria and plasma-membrane NADH oxidoreductase. *Mol. Aspects Med.* **15**, 13-27.
- Lenaz, G., Paolucci, U., Fato, R., D'Aurelio, M., Castelli, G.P., Sgarbi, G., Biagini, G., Ragni, L., Salardi, S. and Cacciari, E. (2002) Enhanced activity of the plasma membrane oxidoreductase in circulating lymphocytes from insulin-dependent diabetes mellitus patients. *Biochem. Biophys. Res. Commun.* **290**, 1589-1592.
- Lisowska, E. (2001) Antigenic properties of human glycophorins – an update. *Adv. Exp. Med. Biol.* **491**, 155-169.
- Liu, Y.C. and Stellwagen, E. (1986) Zonal chromatographic analysis of the interaction of alcohol dehydrogenase with blue-Sepharose. *J. Chromatogr. A* **376**, 149-155.
- Lo, S.C. and Agar, N.S. (1986) NADH-methemoglobin reductase activity in the erythrocytes of newborn and adult mammals. *Experientia* **42**, 1264-1265.
- Löw, H. and Crane, F.L. (1978) Redox function in plasma membranes. *Biochim. Biophys. Acta* **515**, 141-161.
- Löw, H. and Crane, F.L. (1995) The NADH oxidising system of the plasma membrane and metabolic signal control. *Protoplasma* **184**, 158-162.
- Lucchi, L., Bergamini, S., Botti, B., Rapana, R., Ciuffreda, A., Ruggiero, P., Ballestri, M., Tomasi, A. and Albertazzi, A. (2000) Influence of different

hemodialysis membranes on red blood cell susceptibility to oxidative stress. *Artif. Organs* **24**, 1-6.

Lukens, J.N. (1993) Methemoglobinemia and other disorders accompanied by cyanosis. In *Wintrob's Clinical Hematology* (Lee, R.G., Ed.). pp 1262-1271, McGraw Hill, Philadelphia.

Macho, A., Calzado, M.A., Munoz-Blanco, J., Gomez-Diaz, C., Gajate, C., Mollinedo, F., Navas, P. and Munoz, E. (1999) Selective induction of apoptosis by capsaicin in transformed cells: the role of reactive oxygen species and calcium. *Cell Death Differ.* **6**, 155-165.

Mansouri, A. and Winterhalter, K.H. (1974) Nonequivalence of chains in hemoglobin oxidation and oxygen binding. Effect of organic phosphates. *Biochemistry* **13**, 3311-3314.

Marques, F., Crespo, M.E. and Bicho, M. (1995) Control of NADH ferricyanide reductase activity in the human erythrocyte by somatotrophin and insulin. *Redox Rep.* **1**, 113-117.

Marques, F. and Bicho, M.P. (1997) Activation of a NADH dehydrogenase in the human erythrocyte by beta-adrenergic agonists: possible involvement of a G protein in enzyme activation. *Biol. Signals* **6**, 52-61.

Mawatari, S. and Murakami, K. (2001) Effects of ascorbate on membrane phospholipids and tocopherols of intact erythrocytes during peroxidation by t-butylhydroperoxide: comparison with effects of dithiothreitol. *Lipids* **36**, 57-65.

May, J.M., Qu, Z.C. and Whitesell, R.R. (1995a) Ascorbic acid recycling enhances the antioxidant reserve of human erythrocytes. *Biochemistry* **34**, 12721-12728.

May, J.M., Qu, Z.C. and Whitesell, R.R. (1995b) Ascorbate is the major electron donor for a transmembrane oxidoreductase of human erythrocytes. *Biochim. Biophys. Acta* **1238**, 127-136.

May, J.M., Qu, Z.C. and Morrow, J.D. (1996a) Interaction of ascorbate and alpha-tocopherol in resealed human erythrocyte ghosts - transmembrane electron transfer and protection from lipid peroxidation. *J. Biol. Chem.* **271**, 10577-10582.

May, J.M., Qu, Z.C., Whitesell, R.R. and Cobb, C.E. (1996b) Ascorbate recycling in human erythrocytes: role of GSH in reducing dehydroascorbate. *Free Radic. Biol. Med.* **20**, 543-551.

May, J.M., Mendriatta, S., Hill, K.E. and Burk, R.F. (1997) Reduction of dehydroascorbate to ascorbate by the selenoenzyme thioredoxin reductase. *J. Biol. Chem.* **272**, 22601-22610.

May, J.M. (1999) Is ascorbic acid an antioxidant for the plasma membrane? *FASEB J.* **13**, 995-1006.

May, J.M. and Qu, Z.C. (1999) Ascorbate-dependent electron transfer across the human erythrocyte membrane. *Biochim. Biophys. Acta* **1421**, 19-31.

May, J.M., Qu, Z.C. and Cobb, C.E. (2000a) Extracellular reduction of the ascorbate free radical by human erythrocytes. *Biochem. Biophys. Res. Commun.* **267**, 118-123.

May, J.M., Qu, Z.C., Morrow, J.D. and Cobb, C.E. (2000b) Ascorbate-dependent protection of human erythrocytes against oxidant stress generated by extracellular diazobenzene sulfonate. *Biochem. Pharmacol.* **60**, 47-53.

May, J.M., Qu, Z.C. and Cobb, C.E. (2001a) Recycling of the ascorbate free radical by human erythrocyte membranes. *Free Radic. Biol. Med.* **31**, 117-124.

- May, J.M., Qu, Z.C. and Morrow, J.D. (2001b) Mechanisms of ascorbic acid recycling in human erythrocytes. *Biochim. Biophys. Acta* **1528**, 159-166.
- Mazo, A., Gelpí, J.L. and Cortés, A. (1990) Comparison of the kinetic behaviour of lactate dehydrogenase and cytosolic and mitochondrial malate dehydrogenase from guinea pig skeletal muscle. *J. Mol. Catal.* **58**, 269-275.
- Medina, M.A., Sanchez-Jimenez, F., Segura, J.A. and Nunez de Castro, I. (1988) Transmembrane ferricyanide reductase activity in Ehrlich ascites tumor cells. *Biochim. Biophys. Acta* **946**, 1-4.
- Medina, M.A. and Nunez de Castro, I. (1994) Functional reconstitution of Ehrlich cell plasma membrane ferricyanide reductase. *Biochem. Biophys. Res. Commun.* **205**, 1109-1113.
- Meister, A. and Anderson, M.E. (1983) Glutathione. *Annu. Rev. Biochem.* **52**, 711-760.
- Mendiratta, S., Qu, Z.C. and May, J.M. (1998) Enzyme-dependent ascorbate recycling in human erythrocytes: role of thioredoxin reductase. *Free Radic. Biol. Med.* **25**, 221-228.
- Menz, G.L., McCall, M.N. and Kuchel, P.W. (1989) Identification of methyl resonances in the ¹H NMR spectrum of incubated blood cell lysates. *J. Biol. Chem.* **264**, 2100-2107.
- Merker, M.P., Olson, L.E., Bongard, R.D., Patel, M.K., Linehan, J.H. and Dawson, C.A. (1998) Ascorbate-mediated transplasma membrane electron transport in pulmonary arterial endothelial cells. *Am. J. Physiol.* **274**, L685-693.
- Merker, M.P., Bongard, R.D., Kettenhofen, N.J., Okamoto, Y. and Dawson, C.A. (2002) Intracellular redox status affects transplasma membrane electron transport in pulmonary arterial endothelial cells. *Am. J. Physiol.* **282**, L36-43.

- Messana, J.M., Cieslinski, D.A., O'Connor, R.P. and Humes, H.D. (1988) Glutathione protects against exogenous oxidant injury to rabbit renal proximal tubules. *Am. J. Physiol.* **255**, F874-884.
- Miner, C., Lopez-Burillo, S., Garcia-Sancho, J. and Herreros, B. (1983) Plasma membrane NADH dehydrogenase and Ca²⁺-dependent potassium transport in erythrocytes of several animal species. *Biochim. Biophys. Acta* **727**, 266-272.
- Mishra, R.K. and Passow, H. (1969) Induction of intracellular ATP synthesis by extracellular ferricyanide in human red blood cells. *J. Membr. Biol.* **1**, 214-224.
- Mitsuyama, H. and May, J.M. (1999) Uptake and antioxidant effects of ergothioneine in human erythrocytes. *Clin. Sci.* **97**, 407-411.
- Mittler, R. and Zilinskas, B. (1993) Detection of ascorbate peroxidase activity in native gels by inhibition of the ascorbate-dependent reduction of nitroblue tetrazolium. *Anal. Biochem.* **212**, 540-546.
- Moral, A., Lafuente, M.J., Lafuente, A., Castel, T., Balesta, A.M., Trias, M.L. and Group, M.M.M. (2000) The use of erythrocyte glutathione as a predictive marker for malignant melanoma. *Anticancer Res.* **20**, 4757-4760.
- Morel, Y. and Barouki, R. (1999) Repression of gene expression by oxidative stress. *Biochem. J.* **342**, 481-496.
- Morishima, I., Okada, K., Yonezawa, T. and Goto, K. (1971) Carbon-13 contact shifts of a σ -electron system. Conformational dependence of carbon-13 contact shifts in six-membered rings. *J. Am. Chem. Soc.* **93**, 3922 - 3926.
- Morishima, I., Yoshikawa, K. and Okada, K. (1975) Studies on the proton and carbon-13 contact shifts for σ -bonded molecules. Stereospecific electron spin transmission in cyclic and bicyclic amines. *J. Am. Chem. Soc.* **98**, 3787 - 3793.

- Morré, D.J., Morré, D.M. and Wu, L.Y. (1994) Response to adriamycin of transplasma membrane electron transport in adriamycin-resistant and nonresistant HL-60 cells. *J. Bioenerg. Biomembr.* **26**, 137-142.
- Morré, D.J., Chueh, P.J. and Morré, D.M. (1995) Capsaicin inhibits preferentially the NADH oxidase and growth of transformed cells in culture. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 1831-1835.
- Morré, D.J., Caldwell, S., Mayorga, A., Wu, L.Y. and Morré, D.M. (1997) NADH oxidase activity from sera altered by capsaicin is widely distributed among cancer patients. *Arch. Biochem. Biophys.* **342**, 224-230.
- Morré, D.J. (1998) NADH oxidase activity of soybean plasma membranes inhibited by submicromolar concentrations of ATP. *Mol. Cell. Biochem.* **187**, 41-46.
- Morré, D.J., Lawler, J., Wang, S., Keenan, T.W. and Morré, D.M. (2002) Entrainment in solution of an oscillating NADH oxidase activity from the bovine milk fat globule membrane with a temperature-compensated period length suggestive of an ultradian time-keeping (clock) function. *Biochim. Biophys. Acta* **1559**, 10-20.
- Morré, D.J. and Morré, D.M. (2003a) The plasma membrane-associated NADH oxidase (ECTO-NOX) of mouse skin responds to blue light. *J. Photochem. Photobiol. B.* **70**, 7-12.
- Morré, D.J. and Morré, D.M. (2003b) Cell surface NADH oxidases (ECTO-NOX proteins) with roles in cancer, cellular time-keeping, growth, aging and neurodegenerative diseases. *Free Radic. Res.* **37**, 795-808.
- Morré, D.M., Lenaz, G. and Morré, D.J. (2000) Surface oxidase and oxidative stress propagation in aging. *J. Exp. Biol.* **203**, 1513-1521.

Mulquiney, P.J., Bubb, W.A. and Kuchel, P.W. (1999) Model of 2,3-bisphosphoglycerate metabolism in the human erythrocyte based on detailed enzyme kinetic equations: *in vivo* kinetic characterization of 2,3-bisphosphoglycerate synthase/phosphatase using ^{13}C and ^{31}P NMR. *Biochem. J.* **342**, 567-580.

Mulquiney, P.J. and Kuchel, P.W. (1999a) Model of 2,3-bisphosphoglycerate metabolism in the human erythrocyte based on detailed enzyme kinetic equations: equations and parameter refinement. *Biochem. J.* **342**, 581-596.

Mulquiney, P.J. and Kuchel, P.W. (1999b) Model of 2,3-bisphosphoglycerate metabolism in the human erythrocyte based on detailed enzyme kinetic equations: computer simulation and Metabolic Control Analysis. *Biochem. J.* **342**, 597-604.

Nadlinger, K., Westerthaler, W., Storga-Tomic, D. and Birkmayer, J.G. (2002) Extracellular metabolisation of NADH by blood cells correlates with intracellular ATP levels. *Biochim. Biophys. Acta* **1573**, 177-182.

Navas, P., Sun, I.L., Morr , D.J. and Crane, F.L. (1986) Decrease of NADH in HeLa-cells in the presence of transferrin or ferricyanide. *Biochem. Biophys. Res. Commun.* **135**, 110-115.

Navas, P., Estevez, A., Buron, M.I., Villalba, J.M. and Crane, F.L. (1988) Cell surface glycoconjugates control the activity of the NADH-ascorbate free radical reductase of rat liver plasma membrane. *Biochem. Biophys. Res. Commun.* **154**, 1029-1033.

Navas, P., Alcain, F.J., Buron, I., Rodr quez-Aguilera, J.C., Villalba, J.M., Morr , D.M. and Morr , D.J. (1992) Growth factor-stimulated trans plasma membrane electron transport in HL-60 cells. *FEBS Lett.* **299**, 223-226.

Neefjes, V.M., Evelo, C.T., Baars, L.G. and Blanco, C.E. (1999) Erythrocyte glutathione S-transferase as a marker of oxidative stress at birth. *Arch. Dis. Child. Fetal Neonatal Ed.* **81**, F130-133.

- Nijtmans, L.G.J., Henderson, N.S. and Holt, I.J. (2002) Blue Native electrophoresis to study mitochondrial and other protein complexes. *Methods* **26**, 327-334.
- Nisimoto, Y., Wilson, E., Heyl, B.L. and Lambeth, J.D. (1986) NADH dehydrogenase from bovine neutrophil membranes - purification and properties. *J. Biol. Chem.* **261**, 285-290.
- Nisselbaum, J.S., Packer, D.E. and Bodansky, O. (1964) Comparison of the actions of human brain, liver, and heart lactic dehydrogenase variants on nucleotide analogues and on substrate analogues in the absence and in the presence of oxalate and oxamate. *J. Biol. Chem.* **239**, 2830-2834.
- Nualart, F.J., Rivas, C.I., Montecinos, V.P., Godoy, A.S., Guaiquil, V.H., Golde, D.W. and Vera, J.C. (2003) Recycling of vitamin C by a bystander effect. *J. Biol. Chem.* **278**, 10128-10133.
- Oritani, T., Fukuhara, N., Okajima, T., Kitamura, F. and Ohsaka, T. (2004) Electrochemical and spectroscopic studies on electron-transfer reaction between novel water-soluble tetrazolium salts and a superoxide ion. *Inorg. Chim. Acta* **357**, 436-442.
- Orringer, E.P. and Roer, M.E.S. (1979) An ascorbate-mediated transmembrane-reducing system of the human erythrocyte. *J. Clin. Invest.* **63**, 53-58.
- Ottolina, G., Riva, S., Carrea, G., Danieli, B. and Buckmann, A.F. (1989) Enzymatic synthesis of [4R-²H]NAD (P)H and [4S-²H]NAD(P)H and determination of the stereospecificity of 7 alpha- and 12 alpha hydroxysteroid dehydrogenase. *Biochim. Biophys. Acta* **998**, 173-178.
- Papandreou, P. and Rakitzis, E.T. (1989) Determination of NADH₂-ferricyanide oxidoreductase (cytochrome b₅ reductase, diaphorase) activity of human erythrocytes

by an analysis of the time-dependence of NADH₂ oxidation. *Clin. Chim. Acta* **181**, 189-196.

Pendergrass, J.A., Srinivasan, V., Clark, E.P. and Kumar, K.S. (1999) Glutathione redox status in the human cell line, A549, following intracellular glutathione depletion and extracellular glutathione addition. *Tox. Subst. Mech.* **18**, 11-20.

Perego, P., Gatti, L., Carenini, N., Dal Bo, L. and Zunino, F. (2000) Apoptosis induced by extracellular glutathione is mediated by H₂O₂ production and DNA damage. *Int. J. Cancer* **87**, 343-348.

Perrella, M., Shrager, R.I., Ripamonti, M., Manfredi, G., Berger, R.L. and Rossi-Bernardi, L. (1993) Mechanism of the oxidation reaction of deoxyhemoglobin as studied by isolation of the intermediates suggests tertiary structure dependent cooperativity. *Biochemistry* **32**, 5233-5238.

Peskin, A.V. and Winterbourn, C.C. (2000) A microtiter plate assay for superoxide dismutase using a water-soluble tetrazolium salt (WST-1). *Clin. Chim. Acta* **293**, 157-166.

Pretsch, E., Bühlmann, P. and Affolter, C. (2000) *Structure Determination of Organic Compounds. Tables of Spectral Data*. Springer-Verlag, Berlin.

Price, W.S., Roufogalis, B. and Kuchel, P.W. (1989) A simple and inexpensive method for preparing erythrocyte membranes by filtration through a hollow-fibre system. *Anal. Biochem.* **179**, 190-193.

Rabenstein, D.L., Brown, D.W. and McNeil, C.J. (1985) Determination of glutathione in intact and hemolyzed erythrocytes by titration with *tert*-butyl hydroperoxide with end point detection by ¹H nuclear magnetic resonance spectrometry. *Anal. Chem.* **57**, 2294-2299.

Rabenstein, D.L. and Keire, D.A. (1989) Nuclear magnetic resonance spectroscopy of glutathione. In *Coenzymes and Cofactors. Glutathione: Chemical, Biochemical and Medical Aspects* (Dolphin, D., Poulson, R. and Avramovic, O., Ed.). pp 67-101, John Wiley and Sons, New York.

Rae, C.D., Berners-Price, S.J., Bulliman, B.T. and Kuchel, P.W. (1990) Kinetic analysis of the human erythrocyte glyoxalase system using ^1H NMR and a computer model. *Eur. J. Biochem.* **193**, 83-90.

Rae, C.D., Board, P.G. and Kuchel, P.W. (1991) Glyoxalase 2 deficiency in the erythrocytes of a horse: ^1H NMR studies of enzyme kinetics and transport of S-lactoylglutathione. *Arch. Biochem. Biophys.* **291**, 291-299.

Rae, C.D., Sweeney, K., J. E. and Kuchel, P.W. (1993) Stability and nonreactivity of ergothioneine in human erythrocytes studied by ^1H NMR. *Magn. Reson. Med.* **29**, 826-829.

Raftos, J.E., Chapman, B.E., Kuchel, P.W., Lovric, V.A. and Stewart, I.M. (1986) Intra- and extraerythrocyte pH at 37 degrees C and during long term storage at 4 degrees C: ^{31}P NMR measurements and an electrochemical model of the system. *Haematologia.* **19**, 251-268.

Regan, R.F. and Guo, Y. (1999) Extracellular reduced glutathione increases neuronal vulnerability to combined chemical hypoxia and glucose deprivation. *Brain Res.* **817**, 145-150.

Reglinski, J., Hoey, S., Smith, W.E. and Sturrock, R.D. (1988a) Cellular response to oxidative stress at sulfhydryl group receptor sites on the erythrocyte membrane. *J. Biol. Chem.* **263**, 12360-12366.

Reglinski, J., Smith, W.E. and Sturrock, R.D. (1988b) Spin-echo ^1H NMR detected response of ergothioneine to oxidative stress in the intact human erythrocyte. *Magn. Reson. Med.* **6**, 217-223.

Reglinski, J., Smith, W.E., Wilson, R., Buchanan, L.M., McKillop, J.H., Thomsen, J.A., Brzeski, M., Marabani, M. and Sturrock, R.D. (1991) Clinical analysis in intact erythrocytes using ^1H spin echo NMR. *Clin. Chim. Acta* **201**, 45-58.

Renner, E.L., Lake, J.R., Cragoe, E.J., Jr. and Scharschmidt, B.F. (1988) Amiloride and amiloride analogs inhibit Na^+/K^+ -transporting ATPase and Na^+ -coupled alanine transport in rat hepatocytes. *Biochim. Biophys. Acta* **938**, 386-394.

Rodriguez-Aguilera, J.C., Navarro, F., Arroyo, A., Villalba, J.M. and Navas, P. (1993) Transplasma membrane redox system of HL-60 cells is controlled by cAMP. *J. Biol. Chem.* **268**, 26346-26349.

Russell, J., Spickett, C., Reglinski, J., Smith, W., McMurray, J. and Abdullah, I. (1994) Alteration of the erythrocyte glutathione redox balance by *N*-acetylcysteine, captopril and exogenous glutathione. *FEBS Lett.* **347**, 215-220.

Sandelius, A.S., Barr, R., Crane, F.L. and Morr , D.J. (1987) Redox reactions of plasma membranes isolated from soybean hypocotyls by phase partition. *Plant Sci.* **48**, 1-10.

Sandstr m, B.E. and Marklund, S.L. (1990) Effects of variation in glutathione peroxidase activity on DNA damage and cell survival in human cells exposed to hydrogen peroxide and *t*-butyl peroxide. *Biochem. J.* **271**, 17-23.

Sarneski, J.E., Suprenant, H.L., Molen, F.K. and Reilley, C.N. (1975) Chemical shifts and protonation shifts in carbon-13 nuclear magnetic resonance studies of aqueous amines. *Anal. Chem.* **47**, 2116-2124.

Schagger, H. and von Jagow, G. (1991) Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form. *Anal. Biochem.* **199**, 223-231.

- Schipfer, W., Neophytou, B., Trobisch, R., Groiss, O. and Goldenberg, H. (1985) Reduction of extracellular potassium ferricyanide by transmembrane NADH: (acceptor) oxidoreductase of human erythrocytes. *Int. J. Biochem.* **17**, 819-823.
- Shaka, A.J., Keeler, J. and Freeman, R. (1983) Evaluation of a new broad-band decoupling sequence - Waltz-16. *J. Magn. Reson.* **53**, 313-340.
- Shevchenko, A., Wilm, M., Vorm, O. and Mann, M. (1996) Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal. Chem.* **68**, 850-858.
- Simpson, R.J., Brindle, K.M., Brown, F.F., Campbell, I.D. and Foxall, D.L. (1981) Studies of pyruvate-water isotope exchange catalysed by erythrocytes and proteins. *Biochem. J.* **193**, 401-406.
- Simpson, R.J., Brindle, K.M., Brown, F.F., Campbell, I.D. and Foxall, D.L. (1982) Studies of lactate dehydrogenase in the purified state and in intact erythrocytes. *Biochem. J.* **202**, 581-587.
- Slade, N., Storga-Tomic, D., Birkmayer, G.D., Pavelic, K. and Pavelic, J. (1999) Effect of extracellular NADH on human tumor cell proliferation. *Anticancer Res.* **19**, 5355-5360.
- Smith, J.E., Barnes, J.K., Kaneko, J.J. and Freedland, R.A. (1965) Erythrocytic enzymes of various animal species. *Nature* **205**, 298-299.
- Smith, J.E., Kiefer, S. and Lee, M. (1972) Glutathione reduction and other enzyme activities in equine erythrocytes. *Comp. Biochem. Physiol. B.* **43**, 413-417.
- Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K., Gartner, F.H., Provenzano, M.D., Fujimoto, E.K., Goeke, N.M., Olson, B.J. and Klenk, D.C. (1985) Measurement of protein using bicinchoninic acid.[erratum appears in *Anal Biochem* 1987 May 15;163(1):279]. *Anal. Biochem.* **150**, 76-85.

Snyder, L.M., Fortier, N.L., McKenny, J., Trainor, J., Sheerin, H. and Mohandas, N. (1988) The role of membrane plasma sulfhydryl groups in hydrogen peroxide-mediated membrane damage in human erythrocytes. *Biochim. Biophys. Acta* **937**, 229-240.

Spickett, C.M., Reglinski, J., Smith, W.E., Wilson, R., Walker, J.J. and McKillop, J. (1998) Erythrocyte glutathione balance and membrane stability during preeclampsia. *Free Radic. Biol. Med.* **24**, 1049-1055.

Srivastava, S.K., Awasthi, Y.C. and Beutler, E. (1974) Useful agents for the study of glutathione metabolism in erythrocytes. Organic hydroperoxides. *Biochem. J.* **139**, 289-295.

Steck, T.L. and Kant, J.A. (1974) Preparation of impermeable ghosts and inside-out vesicles from human erythrocyte membranes. *Methods Enzymol.* **31**, 172-180.

Stewart, I.M., Chapman, B.E., Kirk, K., Kuchel, P.W., Lovric, V.A. and Raftos, J.E. (1986) Intracellular pH in stored erythrocytes. Refinement and further characterisation of the ³¹P-NMR methylphosphonate procedure. *Biochim. Biophys. Acta* **885**, 23-33.

Stryer, L. (1995) *Biochemistry, 4th edition*. W.H. Freeman and Company, New York.

Sun, I.L., Crane, F.L., Chou, J.Y., Löw, H. and Grebing, C. (1983) Transformed liver cells have modified transplasma membrane redox activity which is sensitive to adriamycin. *Biochem. Biophys. Res. Commun.* **116**, 210-216.

Sun, I.L., Sun, E.E., Crane, F.L., Morré, D.J., Lindgren, A. and Löw, H. (1992) Requirement for coenzyme Q in plasma membrane electron transport. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 11126-11130.

- Sun, I.L., Crane, F.L. and Löw, H. (1994) Bombesin stimulates transplasma-membrane electron transport by Swiss 3T3 cells. *Biochim. Biophys. Acta* **1221**, 206-210.
- Tan, A.S. and Berridge, M.V. (2000) Superoxide produced by activated neutrophils efficiently reduces the tetrazolium salt WST-1 to produce a soluble formazan: a simple colorimetric assay for measuring respiratory burst activation and for screening anti-inflammatory agents. *J. Immunol. Methods* **238**, 59-68.
- Tan, A.S. and Berridge, M.V. (2004) Distinct *trans*-plasma membrane redox pathways reduce cell-impermeable dyes in HeLa cells. *Redox Rep.* in press.
- Tanner, M.J. (1993) The major integral proteins of the human red cell. *Baillieres Clin. Haematol.* **6**, 333-356.
- Tavazzi, B., Di Pierro, D., Amorini, A.M., Fazzina, G., Tuttobene, M., Giardina, B. and Lazzarino, G. (2000) Energy metabolism and lipid peroxidation of human erythrocytes as a function of increased oxidative stress. *Eur. J. Biochem.* **267**, 684-689.
- Tavazzi, B., Amorini, A.M., Fazzina, G., Di Pierro, D., Tuttobene, M., Giardina, B. and Lazzarino, G. (2001) Oxidative stress induces impairment of human erythrocyte energy metabolism through the oxygen radical-mediated direct activation of AMP-deaminase. *J. Biol. Chem.* **276**, 48083-48092.
- Thom, S.R., Kang, M., Fisher, D. and Ischiropoulos, H. (1997) Release of glutathione from erythrocytes and other markers of oxidative stress in carbon monoxide poisoning. *J. Appl. Physiol.* **82**, 1424-1432.
- Thomas, D., Ritz, M.F., Malviya, A.N. and Gaillard, S. (1996) Intracellular acidification mediates the proliferative response of PC12 cells induced by potassium ferricyanide and involves MAP kinase activation. *Int. J. Cancer* **68**, 547-552.

- Thorburn, D.R. and Kuchel, P.W. (1985) Regulation of the human-erythrocyte hexose-monophosphate shunt under conditions of oxidative stress. A study using NMR spectroscopy, a kinetic isotope effect, a reconstituted system and computer simulation. *Eur. J. Biochem.* **150**, 371-386.
- Tietze, F. (1969) Enzymic method for quantitative determination of nanogram amounts of total and oxidized glutathione: applications to mammalian blood and other tissues. *Anal. Biochem.* **27**, 502-522.
- Tomoda, A., Takeshita, M. and Yoneyama, Y. (1978) Analysis of met-form hemoglobin in glucose-deleted human red cells. *FEBS Lett.* **88**, 247-250.
- Turner, C.J., Connolly, P.J. and Stern, A.S. (1999) Artifacts in sensitivity-enhanced HSQC. *J. Magn. Reson.* **137**, 281-284.
- Ukeda, H., Kawana, D., Maeda, S. and Sawamura, M. (1999) Spectrophotometric assay for superoxide dismutase based on the reduction of highly water-soluble tetrazolium salts by xanthine-xanthine oxidase. *Biosci. Biotechnol. Biochem.* **63**, 485-488.
- Vaillant, F., Larm, J.A., McMullen, G.L., Wolvetang, E.J. and Lawen, A. (1996) Effectors of the mammalian plasma membrane NADH-oxidoreductase system. Short-chain ubiquinone analogues as potent stimulators. *J. Bioenerg. Biomembr.* **28**, 531-540.
- Van Duijn, M.M., Vanderzee, J., Vansteveninck, J. and Vandenbroek, P.J.A. (1998) Ascorbate stimulates ferricyanide reduction in HL-60 Cells through a mechanism distinct from the NADH-dependent plasma membrane reductase. *J. Biol. Chem.* **273**, 13415-13420.
- Van Duijn, M.M., Tijssen, K., VanSteveninck, J., Van der Broek, P. and Van der Zee, J. (2000) Erythrocytes reduce extracellular ascorbate free radicals using intracellular ascorbate as an electron donor. *J. Biol. Chem.* **275**, 27720-27725.

- Van Duijn, M.M., Buijs, J.T., Van der Zee, J. and Van den Broek, P.J. (2001a) The ascorbate: ascorbate free radical oxidoreductase from the erythrocyte membrane is not cytochrome *b*₅₆₁. *Protoplasma* **217**, 94-100.
- Van Duijn, M.M., Van der Zee, J. and Van den Broek, P.J. (2001b) The ascorbate-driven reduction of extracellular ascorbate free radical by the erythrocyte is an electrogenic process. *FEBS Lett.* **491**, 67-70.
- Van Gorp, R.M., Broers, J.L., Reutelingsperger, C.P., Bronnenberg, N.M., Hornstra, G., Van Dam-Mieras, M.C. and Heemskerk, J.W. (1999) Peroxide-induced membrane blebbing in endothelial cells associated with glutathione oxidation but not apoptosis. *Am. J. Physiol.* **277**, C20-28.
- Van Renswoude, J. and Kempf, C. (1984) Purification of integral membrane proteins. *Methods Enzymol.* **104**, 329-339.
- Vijaya, S., Crane, F.L. and Ramasarma, T. (1984) A vanadate-stimulated NADH oxidase in erythrocyte membrane generates hydrogen peroxide. *Mol. Cell. Biochem.* **62**, 175-185.
- Villalba, J.M., Navarro, F., Cordoba, F., Serrano, A., Arroyo, A., Crane, F.L. and Navas, P. (1995) Coenzyme Q reductase from liver plasma membrane: purification and role in trans-plasma-membrane electron transport. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 4887-4891.
- Voegtlin, C., Johnson, J.M. and Dyer, H.A. (1925) Quantitative estimation of the reducing power of normal and cancer tissue. *J. Pharmacol. Exp. Ther.* **24**, 305-334.
- Wang, C.S. and Alaupovic, P. (1978) Isolation and partial characterization of human erythrocyte membrane NADH: (acceptor) oxidoreductase. *J. Supramol. Struct.* **9**, 1-14.

Washburn, M.P. and Wells, W.W. (1999) Identification of the dehydroascorbate acid reductase and thiotransferase (glutaredoxin) activities of bovine erythrocyte glutathione peroxidase. *Biochem. Biophys. Res. Commun.* **257**, 567-571.

Wells, W.W., Xu, D.P. and Washburn, M.P. (1995) Glutathione: dehydroascorbate oxidoreductases. *Methods Enzymol.* **252**, 30-38.

Willis, J. and Schleich, T. (1996) Oxidative-stress induced protein glutathione mixed-disulfide formation in the ocular lens. *Biochim. Biophys. Acta* **1313**, 20-28.

Wolvetang, E.J., Larm, J.A., Moutsoulas, P. and Lawen, A. (1996) Apoptosis induced by inhibitors of the plasma membrane NADH-oxidase involves Bcl-2 and calcineurin. *Cell Growth Differ.* **7**, 1315-1325.

Yamamoto, S.I. and Kawasaki, T. (1981) The involvement of the membrane oxidoreduction system in stimulating amino-acid-uptake in Ehrlich ascites tumor-cells. *Biochim. Biophys. Acta* **644**, 192-200.

Ye, B., Gitler, C. and Gressel, J. (1997) A high-sensitivity, single-gel, polyacrylamide gel electrophoresis method for the quantitative determination of glutathione reductases. *Anal. Biochem.* **246**, 159-165.

York, M.J., Kuchel, P.W. and Chapman, B.E. (1984) A proton nuclear magnetic resonance study of gamma-glutamyl-amino acid cyclotransferase in human erythrocytes. *J. Biol. Chem.* **259**, 15085-15088.

Yucel, G., Gumuslu, S., Yesilkaya, A., Akin, H. and Aksu, A. (2000) The chemiluminescence detection of protective effect of extracellularly added NADPH and NADH on erythrocytes against cumene hydroperoxide-induced oxidative stress. *Turk. J. Med. Sci.* **26**, 187-191.

- Zamudio, I. and Canessa, M. (1966) Nicotinamide-adenine dinucleotide dehydrogenase activity of human erythrocyte membranes. *Biochim. Biophys. Acta* **120**, 165-169.
- Zamudio, I., Cellino, M. and Canessa-Fischer, M. (1969) The relation between membrane structure and NADH: (acceptor) oxidoreductase activity of erythrocyte ghosts. *Arch. Biochem. Biophys.* **129**, 336-345.
- Zavodnik, L.B., Zavodnik, I.B., Niekurzak, A., Szosland, K. and Bryszewska, M. (1998) Activation of red blood cell glutathione peroxidase and morphological transformation of erythrocytes under the action of *tert*-butyl hydroperoxide. *Biochem. Mol. Biol. Int.* **44**, 577-588.
- Zemkova, H., Teisinger, J. and Vyskocil, F. (1984) Hyperpolarisation of mouse skeletal muscle plasma membrane induced by extracellular NADH. *Biochim. Biophys. Acta* **775**, 64-70.
- Zerez, C.R., Lachant, N.A. and Tanaka, K.R. (1990) Impaired erythrocyte methemoglobin reduction in sickle cell disease: dependence of methemoglobin reduction on reduced nicotinamide adenine dinucleotide content. *Blood* **76**, 1008-1014.
- Zinkl, J. and Kaneko, J.J. (1973) Erythrocytic enzymes and glycolytic intermediates in the normal bovine and in bovine erythropoietic porphyria. *Comp. Biochem. Physiol. A.* **45**, 463-476.
- Zou, C.G., Agar, N.S. and Jones, G.L. (2000) Haemolysis of human and sheep red blood cells in glycerol media: the effect of pH and the role of band 3. *Comp. Biochem. Physiol., A: Mol. Integrative Physiol.* **127**, 347-353.
- Zou, C.G., Agar, N.S. and Jones, G.L. (2001) Oxidative insult in sheep red blood cells induced by *t*-butyl hydroperoxide: The roles of glutathione and glutathione peroxidase. *Free Radic. Res.* **34**,

Zurbriggen, R. and Dreyer, J.-L. (1994) An NADH-diaphorase is located at the cell plasma membrane in a mouse neuroblastoma cell line NB41A3. *Biochim. Biophys. Acta* **1183**, 513-520.

Zurbriggen, R. and Dreyer, J.-L. (1996) The plasma membrane NADH-diaphorase is active during selective phases of the cell cycle in mouse neuroblastoma cell line NB41A3. Its relation to cell growth and differentiation. *Biochim. Biophys. Acta* **1312**, 215-222.