

Speciation and phytoavailability of arsenic in contaminated soils using a SEP and XANES spectroscopy³

ABSTRACT

In this study, a sequential extraction procedure (SEP) and X-ray absorption near edge structure (XANES) spectroscopy were used to determine the solid-phase speciation and phytoavailability of arsenic (As) of historically contaminated soils from As containing pesticides and herbicides and soils spiked with As in the laboratory. *Brassica juncea* was grown in the contaminated soils to measure plant available As in a glasshouse experiment. Arsenic associated with amorphous Fe oxides was found to be the dominant phase using both SEP and XANES spectroscopy. Arsenic predominantly existed in arsenate (As^{V}) form in the soils; in few samples As was also present in arsenite (As^{III}) form or in scorodite mineral. Arsenic concentration in shoots showed significant ($P < 0.001$ – 0.05) correlations with the exchangeable As ($r = 0.85$), and amorphous Fe oxides associated As evaluated by the SEP ($r = 0.67$); and As associated with amorphous Fe oxides as determined by XANES spectroscopy ($r = 0.51$). The results show that As in both fractions is readily available for plant uptake and may pose potential risk to the environment. The combination of SEP and XANES spectroscopy allowed the quantitative speciation of As in the contaminated soils and the identification of valence and mineral forms of As. Such detailed knowledge on As speciation and availability is vital for management and rehabilitation of As-contaminated soils.

³ This chapter has been published in *Environmental Science & Technology*, 45 (17): 7135–7142, under the title 'Arsenic speciation and phytoavailability in contaminated soils using a sequential extraction procedure and XANES spectroscopy'. Authors are Nabeel K. Niazi, Balwant Singh and Pushan Shah.

7.1 INTRODUCTION

Arsenic contamination of soil is a major environmental concern due to the toxic and carcinogenic nature of some As compounds (Smith et al. 1998; Mandal and Suzuki 2002). Agricultural use of As-based pesticides and herbicides, ore mining and smelting, CCA treatment of wood and pesticides manufacturing processes are the major anthropogenic sources of soil As contamination (Smith et al. 1998; Mandal and Suzuki 2002). Arsenicals (as sodium arsenite) were applied to control the southward migration of cattle tick across New South Wales (NSW) in Australia from early 1900s to 1955. Over 1500 dips were constructed along the eastern coast of Australia with many of these are located in northern NSW (Smith et al. 1998). Arsenic-containing herbicides were also used to suppress weed growth along the former railway corridors in South Australia over 30 years ago (Smith et al. 2006a). The repeated application of As-based pesticides and herbicides led to As concentrations of up to 3,000 and 1,400 mg kg⁻¹ in soils at the cattle-dip sites and railway corridors, respectively (Smith et al. 1998; Smith et al. 2006a). Arsenic in the soils adjoining these sites significantly exceeds the ecological investigation level (EIL, 20 mg kg⁻¹) in the soil in Australia (NEPM 1999). Limited research suggests that As exists in more soluble and toxic forms at these sites and thus present significant environmental and health risks (McLaren et al. 1998; Smith et al. 1998). The site-specific evaluation of As-contaminated soil is, therefore, imperative to determine the chemical forms of As and to evaluate phytoavailability of As. Such information would be important for risk assessment and for the development and implementation of suitable management and remediation strategies.

Sequential extraction procedures have been previously used to delineate chemical forms of As in soils such as non-specifically and specifically sorbed As, As occluded in amorphous and

crystalline Fe oxides, and recalcitrant As (McLaren et al. 1998; Lombi et al. 2000; Wenzel et al. 2001; Novoa-Munoz et al. 2007). In a previous study, McLaren et al. (1998) used a sequential fractionation method for As partitioning in contaminated soils. They reported that As was mostly adsorbed onto amorphous Fe oxides (0.1 M NaOH extractable) in soils sampled from disused cattle-dip sites, and a substantial proportion of As was highly labile (0.05 M NaHCO₃ extractable). The sequential extraction procedures are reported to have some limitations which include the partial dissolution of a required fraction (La Force and Fendorf 2000), dissolution of the nontarget phase and a partial recovery of the desired phase due to readsorption or reprecipitation reactions (Ostergren et al. 1999; Wenzel et al. 2001; Scheinost et al. 2002). The redox forms of As in soils are also not detected using a sequential extraction procedure (Wenzel et al. 2001; Novoa-Munoz et al. 2007).

X-ray absorption fine structure (XAFS) spectroscopy is a powerful technique, which has been used to determine the oxidation state and chemically bound forms of As in contaminated soils (Paktunc et al. 2003; Cances et al. 2005; Arai et al. 2006; Meunier et al. 2010). Cances et al. (2005) observed that As predominantly occurred as As^V and approximately 80% of the As was bound to the amorphous Fe oxides throughout the soil profile at a former arsenical processing plant. Arai et al. (2006) investigated an As-contaminated site close to a pesticide (PbHAsO₄) manufacturing plant, and observed that As in the oxidized (0–10 cm) layer was mainly present as As^V in association with the amorphous Fe oxides (71%). Recently, Meunier et al. (2010) determined mineralogical composition of As in As-enriched tailings and soils from abandoned gold mine sites using XANES spectroscopy. They also studied the effect of various mineral forms of As on the bioaccessibility of As using an *in vitro* physiologically based extraction test (PBET). The authors found that As was present in mineral forms such as

arsenopyrite, scorodite, realgar and kankite in the studied samples. The bioaccessibility of As was < 1% of the total As in arsenopyrite and scorodite, and ranged from 2–7% in As-bearing amorphous Fe arsenate and Fe oxyhydroxides minerals (Meunier et al. 2010).

Previous research, however, has not used combination of SEP and XANES spectroscopy, which could be very powerful in delineating various chemical (labile, sorbed, mineral bound), mineral and oxidation states of As in contaminated soils. Therefore, this study was conducted employing SEP and XANES spectroscopy to determine As forms. In addition, the phytoavailability of As in the soils was estimated in a glasshouse experiment to indicate the bioavailability of As in terms of plant available As in a range of As-contaminated soils. The primary objectives of this study were to (1) determine the solid-state speciation of As in contaminated soils sampled from historical cattle-dip sites and railway corridor, and laboratory spiked soils using the combination of a SEP and XANES spectroscopy, and (2) evaluate the phytoavailability of As in soils using *Brassica juncea* as a test plant.

7.2 MATERIALS AND METHODOLOGY

7.2.1 Soil sampling

Surface (0–10 cm) samples were collected from 16 As-contaminated cattle-dip sites in northern NSW and along 3 railway corridors in South Australia, using a (Ø 6 cm) stainless steel soil corer. Sub-surface samples at a depth of 30–40 cm were also taken from Blackwoods and Sandyarm sites, hereafter referred to as Blackwoods30–40 cm and Sandyarm30–40 cm. A number of soil cores (8–10) were collected from each of the contaminated sites, which were mixed together to make a composite soil sample (1.5–2 kg)

for the site. To obtain a wide range of As-contaminated soils for the study, four soils, Westwood, Lansdowne, Lucerne and Narrabri, with a wide ranging pH, free Fe and clay contents, were spiked with As (Figure 7.1; Table 7.1). Except for the sandy Lansdowne soil, the soils were spiked with three levels of As (100, 200 and 400 mg kg⁻¹ As using Na₂HAsO₄·7H₂O); the Lansdowne soil was spiked with 10 and 20 mg kg⁻¹ As.




Figure 7.1 Locations of the As-contaminated cattle-dip sites, railway corridor and As spiked soil samples collected from New South Wales (NSW) and South Australia (SA) states of Australia.

7.2.2 Physicochemical analysis

The soil samples were dried at 40 °C, ground to obtain < 2 mm fraction and homogenised. These samples were used for the glasshouse experiment, and for the laboratory and spectroscopic analyses.

The concentration of Fe and Al in the DCB extract and Fe in ammonium oxalate extract was analysed using an ICP-AES. The relative standard deviation was below 2% for Fe and Al analysis. Organic carbon in the soil samples was determined using CNS analyser (Elementar®). The selected physico-chemical properties of the soil samples are presented in Table 7.1 (see Appendix 4). For the total As analysis in soils, sub-samples (approximately 15 g) were ground to < 200 µm and digested in acids using the methodology described by Huang and Fujii (1996) (Section 4.2.4.1, Chapter 4).

7.2.3 Phytoavailability pot experiment

Small size plastic pots (Ø 8 cm; 0.5 kg capacity) internally lined with the polythene bags were filled with the As-contaminated (cattle-dip sites and railway corridor) and spiked soils (350 g in each pot). The contaminated and spiked soils were irrigated with deionised water and fertilised with basic nutrients including N-P-K and S applied @ 100-40-50 and 25 mg kg⁻¹, respectively. After 3–4 days incubation time, *Brassica juncea* L. (Broad leaf cultivar; Indian mustard) seeds were sown in the moist soil just below the soil surface (5–8 mm deep). After one week of germination, four uniform plants were maintained per pot. The plants were irrigated daily with deionised water to keep the soil moist at the water holding capacity.

Table 7.1 Physico-chemical properties of soil samples from historically As-contaminated cattle-dip and railway corridor sites and soils spiked with As in the laboratory.

Soil samples	pH (1:5, 0.01 M CaCl ₂)	Organic carbon (%)	Clay (%)	CEC (mmol _c kg ⁻¹)	DCB Fe ^a (wt %)	DCB Al ^a (wt %)	Amorphous Fe ^b (wt %)	Crystalline Fe ^b (wt %)
<i>Cattle-dip sites (n = 18)</i>								
Dungarubba	4.75	3.92	6	77	0.22	0.06	0.21	0.01
Hendersons	5.58	4.31	16	90	1.22	0.14	0.79	0.43
Piora	5.93	3.12	32	91	1.33	0.13	0.53	0.81
Sneesbyslane	4.8	5.27	24	91	1.30	0.14	0.46	0.84
JBM	6.01	4.69	20	91	1.88	0.23	1.16	0.72
Boorie	5.74	3.63	32	91	1.34	0.12	0.92	0.42
Henrywhite	5.49	4.24	40	91	1.64	0.18	0.86	0.78
Enak	4.15	9.98	20	89	1.37	0.26	0.59	0.78
Faulkners	4.48	2.81	48	88	6.10	0.85	0.67	5.40
Dunoon	5.37	5.19	40	91	5.47	0.78	0.69	4.78
Nimbin	5.47	1.44	56	91	1.80	0.23	0.44	1.36
Blackwoods	5.43	13.88	10	88	0.72	0.20	0.39	0.32
Blackwoods30–40 cm	5.17	6.82	10	87	0.97	0.26	0.53	0.44
Alexanders	5.80	6.62	24	89	1.94	0.22	0.63	1.31

Table 7.1 Continued...

Soil samples	pH (1:5, 0.01 M CaCl ₂)	Organic carbon (%)	Clay (%)	CEC (mmol _c kg ⁻¹)	DCB Fe ^a (wt %)	DCB Al ^a (wt %)	Amorphous Fe ^b (wt %)	Crystalline Fe ^c (wt %)
Goolmangar	7.02	2.06	44	91	1.12	0.11	0.68	0.44
Sandyarm	5.66	6.04	4	82	0.07	0.02	0.07	0.002
Sandyarm30–40 cm	5.33	1.89	4	58	0.080	0.01	0.078	0.003
Wollongbar	4.82	4.52	44	88	15.9	1.27	0.80	15.10
<i>Railway corridor (n = 3)</i>								
Murray Bridge	6.93	1.84	4	31	0.52	0.05	0.07	0.45
Adelaide hills	5.13	3.42	12	47	1.64	0.17	0.14	1.50
Kapunda	6.70	6.24	23	47	2.57	0.14	0.43	2.14
<i>As spiked soils (n = 4)</i>								
Lansdowne	5.34	1.49	8	48.3	0.49	0.11	0.12	0.38
Westwood	5.42	1.93	20	46	3.30	0.22	0.46	2.83
Narrabri	7.53	0.64	60	234	1.28	0.29	0.23	1.10
Lucerne	4.69	3.47	20	46	1.87	0.17	0.24	1.63

^aDithionite citrate-bicarbonate extractable free Fe and Al; ^bNH₄ oxalate-oxalic acid extractable Fe, 4 hour shaking in the dark at room temperature; ^ccrystalline Fe phase was calculated by subtracting the amorphous Fe concentration from the free Fe concentration.

For each soil three replicates were used in the experiment. *B. juncea* was grown in the glasshouse for eight weeks; the temperature in the glasshouse ranged from 16 to 30°C during the experimental period. After eight weeks, shoots were harvested. The harvested material was washed using a three step washing procedure (tap water-0.1 M HCl-deionised water) as described before (Section 4.2.4, Chapter 4), and dried at 70 °C for 48 hours. The shoot dry matter yield was recorded, the samples were then ground (< 1 mm) and digested in a mixture (1:1) of concentrated nitric and perchloric acids (Miller 1998) for As analysis (Appendices 4 and 5).

7.2.4 Sequential extraction procedure

A four step slightly modified SEP of Wenzel et al. (2001) was employed to partition As in the soil samples. In the modified SEP, Step-I was repeated twice and a washing step using 0.05 M NaCl solution was included after each extraction step (Table 7.2). The washing solution was then mixed with the preceding step solution. The washing step was introduced to remove As in the entrained solution after each extraction step and thus minimise overlap between steps and readsorption or precipitation of As. Step-I involved the extraction of exchangeable or readily available As (0.05 M $(\text{NH}_4)_2\text{SO}_4$), and Step-II released specifically sorbed As fraction using phosphate solution (0.05 M $\text{NH}_4\text{H}_2\text{PO}_4$). The third step involved the extraction of As associated with amorphous Fe oxides (0.2 M $(\text{NH}_4)_2\text{C}_2\text{O}_4/\text{H}_2\text{C}_2\text{O}_4$), and Step-IV dissolved As bound to crystalline Fe oxides (0.2 M $(\text{NH}_4)_2\text{C}_2\text{O}_4/\text{H}_2\text{C}_2\text{O}_4$ + 0.1 M $\text{C}_6\text{H}_8\text{O}_6$). The samples were centrifuged at 3000 rpm for 15 min after each extraction step, filtered using a Whatman[®] no. 1 filter paper and acidified by adding 1 M HCl before As analysis. Arsenic in the residual phase was calculated by subtracting the sum of the four fractions of the SEP from the total As concentration in soil.

Arsenic in soil and plant digests and in soil extracts was determined using a Varian[®] 220Z HG-AAS spectrometer. The QA/QC details are given in Appendix 4.

Table 7.2 Detail for the extraction steps used in the sequential extraction procedure employed for the As speciation of As-contaminated cattle-dip sites, railway corridor and As spiked soils.

Step*	Extractant	Soil:solution ratio	Conditions	Extractable phase
I	0.05 M (NH ₄) ₂ SO ₄	1:30	shaking twice for 2 h at room temperature	exchangeable
II	0.05 M NH ₄ H ₂ PO ₄	1:25	shaking for 12 h at room temperature	specifically sorbed
III	0.2 M (NH ₄) ₂ C ₂ O ₄ /H ₂ C ₂ O ₄ , pH 3.0	1:25	shaking for 4 h in dark at room temperature	bound to amorphous Fe oxides
IV	0.2 M (NH ₄) ₂ C ₂ O ₄ /H ₂ C ₂ O ₄ + 0.1 M C ₆ H ₈ O ₆ , pH 3.0	1:25	heating in water bath in light for 30 min at 96±3 °C	bound to crystalline Fe oxides

* A washing step using 10 mL of 0.05 M NaCl solution was included after each extraction step

7.2.5 XANES spectroscopy

The reference compounds used for the XANES spectroscopy included scorodite (FeAsO₄·2H₂O), As^V adsorbed to goethite (hereafter referred to as As^V-Gt), As^{V/III} adsorbed

to ferrihydrite ($\text{As}^{\text{V/III}}\text{-Fh}$) (Grafe et al. 2004), As^{V} coprecipitated with ferrihydrite ($\text{As}^{\text{V}}\text{-Cop-Fh}$) (Voegelin et al. 2007), prepared following the methods described in detail in Appendix 4. Natural samples of arsenopyrite (FeAsS), orpiment (As_2S_3) and realgar (AsS) obtained from South Australian Museum were also included as reference compounds. Two scans in the XANES region were collected for each of the reference compound and soil samples. The experimental details for the XANES analysis are given Appendix 4. XANES spectra for two samples (Boorie and Wollongbar) were also obtained after Step-II and Step-III of the SEP.

The oxidation state of As was found to be associated with a shift in the white line energy position of As^{III} (11,872.2 eV) and As^{V} (11,875.7 eV) (Kachenko et al. 2010). The software, *Average* was used for averaging the raw spectra, while smoothing, normalisation and linear least-squares combination fits (LCF) were performed using *Athena* (Ravel and Newville 2005; Shah et al. 2007). LCF of the XANES spectra of reference compounds with those of the soil samples was employed to estimate As forms in the soil.

Linear combination fitting using *Athena*

Athena is a XAFS graphical utility software used for the background removal, smoothing and normalisation of the XANES data (Ravel and Newville 2005; Shah et al. 2007). Linear least-square combination fitting (LCF) of XANES spectra of reference compounds to those of the unknown soil samples was performed using a combination of mathematical functions such as the Gaussian, Lorentzian and arctangent step functions, in *Athena* (Shah et al. 2007). The prerequisite for successful LCF analysis is that the data should be correctly normalised. The software allowed the option for selecting up to four reference compounds to build the fit for an unknown sample. It allowed restricting the weight of reference compounds between 0 and

1. The R-factor value was calculated to determine the quality of the fit. The software presented results with different combinations (of reference compounds) with the corresponding R-factor values calculated for each combination. The best combination was selected based on the criteria that described the goodness of the fit obtained for the spectra of an unknown sample in terms of the peak height, the post-edge region, and a minimum R-factor (i.e., normalised sum of residuals) value (Manning 2005; Ravel and Newville 2005) that is equivalent to the normalised sum of the residuals:

$$\text{Normalised sum of residuals (R-factor)} = \sum ((y - y_{fit})^2) / \sum (y^2)$$

In addition, the known chemistry of the soils was also taken into account that appeared to be helpful in selecting the reference compounds used for the linear combination analysis, in the studied samples.

Using the aforementioned criteria, 4 reference compounds, namely As^V-Fh, As^V-Gt, As^{III}-Fh and scorodite were used for the LCF of XANES spectra of all soil samples with the exception of Sandyarm and Sandyarm30–40 cm samples (see next in Table 7.4), where the best fit was obtained with a reduced R-factor value by replacing scorodite with orpiment; in the case of Piora and Murray Bridge samples, the inclusion of As^V-Cop-Fh instead of scorodite, yielded the best fit.

7.2.6 Data analysis

The correlations and basic statistical analysis were performed using GenStat version 12.1 (Payne et al. 2009). The amounts of As (mg kg⁻¹) associated with amorphous and crystalline Fe oxides phases in soil were calculated from the LCF predicted proportions of these fractions using XANES spectroscopy.

7.3 RESULTS AND DISCUSSION

Soil pH for majority of the samples was < 6 except for the Goolmangar (7.02), Murray Bridge (6.93), Adelaide Hills (6.70) and Narrabri (7.53) soils (Table 7.1). Free Fe concentration varied between 0.07 and 15.9 wt % in the soils with < 1 wt % in the lighter textured soils i.e., Dungarubba, Blackwoods, Blackwoods30–40 cm, Sandyarm, Sandyarm30–40 cm, Murray Bridge and Lansdowne samples. Three samples (Faulkners, Dunoon, Wollongbar) contained > 5 wt % free Fe, with a notably high Fe concentration (15.9 wt %) in the Wollongbar soil. Amorphous Fe concentration varied between 0.07 and 1.16 wt % in the soil samples (Table 7.1) and it constituted 5.06–97.9% of the free Fe content with a mean value of 40.1%.




Figure 7.2 Sequential extraction procedure applied to the cattle-dip sites, railway corridor and As spiked soil samples ($n = 18, 3$ and 11 , respectively). Means of each extracted fraction are presented as percent (%) of the total soil As concentration present in the contaminated and As spiked soils. The residual fraction of As was calculated by subtracting sum of the four fractions from the total As concentration in each of the soils. The bars represent \pm SD around the mean values.

7.3.1 Total As in soil

Total As concentration in the soils ranged between 10 and 1,406 mg kg⁻¹ (Table 7.3); for cattle-dip sites, As ranged from 21–1,406 mg kg⁻¹ whereas for the railway corridor soils from 119–257 mg kg⁻¹. Dungarubba, Sandyarm and Sandyarm30–40 cm soils contained relatively low As concentration with values ≤ 34 mg kg⁻¹. Total As concentration increased with depth for the two samples (Sandyarm and Blackwoods), which indicated possible leaching of As in these sandy soils, which is in agreement with the previous results (Kimber et al. 2002).

7.3.2 Arsenic fractionation in soils using a SEP

Exchangeable (readily available) As (Step-I) constituted less than 3.5% of the total soil As, except for three As spiked soils, Lansdowne, Narrabri and Lucerne where it was significantly higher, with values up to 34.7% (Table 7.3; Figure 7.2). The greater amount of As released during the first extraction step in As spiked soils (see Figure 7.2) could be related to the soluble form of added As and shorter equilibration time of added As (6 weeks) in the spiked soils (Smith et al. 1998; Tang et al. 2007; Juhasz et al. 2008). Negligible As removal (< 0.3% of the total soil As) in the exchangeable pool occurred in Faulkners, Dunoon and Wollongbar samples, despite of their high total As contents (> 400 mg kg⁻¹), which can be attributed to the presence of high free Fe contents (> 5 wt %) in these samples. Conversely, relatively higher proportion of As (3.1%) was extracted in the Sandyarm sample, which could be due to the lowest free Fe concentration (0.07 wt %) in this sample.

Table 7.3 Arsenic distribution in various phases as determined by the SEP showing the As distribution in cattle-dip sites, railway corridor and As spiked soil samples ($n = 32$).

Soil samples	% extractable As in each step of SEP ^a				Total soil As ^b	As in shoot ^a
	Step-I	Step-II	Step-III	Step-IV	(mg kg ⁻¹)	(mg kg ⁻¹ DW)
<i>Cattle-dip sites (n = 18)</i>						
Dungarubba	0.3±0.05	8.0±0.4	67±9.0	10±0.9	21±1	3.8±0.12
Hendersons	1.3±0.16	8.0±0.7	75±2.1	15±0.1	95±2	3.6±0.8
Piora	0.9±0.06	16±1.6	56±1.6	15±1.3	253±4	7.9±0.4
Sneesbylane	0.6±0.06	14±1.0	55±3.1	15±0.8	209±17	8.1±0.7
JBM	1.9±0.12	21±0.7	53±1.0	16±0.9	799±138	7.0±0.5
Boorie	1.3±0.03	18±0.6	49±0.8	10±0.5	1406±61	31.1±2.4
Henrywhite	0.4±0.02	16±0.3	59±3.8	17±1.3	643±64	12.7±1.2
Enak	1.1±0.03	6±0.2	61±0.8	13±0.6	778±156	21.7±1.7
Faulkners	0.1±0.01	14±0.7	42±1.7	38±2.2	423±4	4.8±1.2
Dunoon	0.2±0.01	13±0.9	43±0.1	41±0.2	583±15	6.3±0.9
Nimbin	0.2±0.04	17±1.4	49±1.8	24±2.4	84±1	3.7±0.2
Blackwoods	1.1±0.04	6±0.4	59±4.2	10±1.7	249±22	7.2±0.4
Blackwoods30–40 cm	3.3±0.17	8±0.3	71±6.6	9±0.1	782±101	74.8±2.1
Alexanders	1.5±0.05	18±0.4	46±1.7	18±1.2	548±10	14.6±1.9
Goolmangar	0.5±0.03	24±1.0	62±2.3	12±0.1	174±6	4.4±0.4
Sandyarm	3.1±0.37	11±0.6	57±5.6	8±0.1	29±1	6.9±1.2
Sandyarm30–40 cm	1.9±0.04	9±1.8	59±6.6	9±0.6	34±3	8.1±2.2
Wollongbar	0.1±0.01	10±0.4	49±2.3	38±1.4	481±5	4.2±0.5
<i>Railway corridor (n = 3)</i>						
Murray Bridge	2.5±0.16	20±1.4	50±1.8	17±1.5	119±6	14.4±1.7
Adelaide hills	1.2±0.17	9±0.2	53±0.9	34±1.7	257±14	13.9±3.8
Kapunda	1.1±0.02	8±0.8	69±6.7	15±2.1	211±9	10.3±0.9
<i>As spiked soils (n = 11)</i>						
Lansdowne-10	28.4±0.24	31±0.9	39±0.6	7±0.9	11±0.3	10.1±0.8
Lansdowne-20	34.7±0.48	43±3.4	34±0.8	5±0.1	23±1	44.5±3.5
Westwood-100	2.0±0.12	29±0.7	52±5.6	19±1.0	102±5	6.2±1.1

Table 7.3. Continued...

Soil samples	% extractable As in each step of SEP				Total soil As	As in shoot
	Step-I	Step-II	Step-III	Step-IV	(mg kg ⁻¹)	(mg kg ⁻¹ DW)
Westwood-200	1.9±0.13	37±0.8	51±1.7	11±1.1	203±7	12.4±1.2
Westwood-400	2.2±0.10	27±1.4	57±2.1	13±1.3	404±19	12.1±1.3
Narrabri-100	6.4±0.20	66±3.5	22±1.2	6±0.3	103±3	11.7±0.8
Narrabri-200	11.8±0.81	54±2.1	29±1.8	6±0.2	205±4	20.0±1.2
Narrabri-400	16.1±0.38	52±1.6	28±1.2	4±0.2	405±9	mv
Lucerne-100	10.7±1.69	37±2.0	49±1.9	7±0.4	104±4	16.1±0.5
Lucerne-200	15.7±1.71	33±1.5	43±2.0	9±0.8	202±6	22.3±0.3
Lucerne-400	10.3±0.26	40±0.5	44±0.3	7±0.6	405±11	mv

mv = missing value (plants died after germination); the results are presented as % mean values ± SD; ^a*n* = 3; ^b*n* = 2

The specifically sorbed As (Step-II) was the second dominant phase, and it constituted 6–66% of the total As. Similar to the exchangeable As phase, the As spiked soils showed a greater proportion of As in this fraction with maximum As extraction in the Narrabri soil (66%) spiked with 100 mg kg⁻¹ As (Table 7.3).

The substantial release of As in Step-II indicated that As can be potentially mobilised as a result of phosphate application to these soils (Smith et al. 1998; Tao et al. 2006). However, further studies are warranted to determine the effect of phosphate on the mobility and bioavailability of As to plants under field conditions. For all samples, As was primarily associated with amorphous Fe oxides (Step-III) and it ranged from 22–75% with a mean of 51% of the total As. Relatively, lower values were obtained in the Step-III for the As spiked soils, as higher proportion of As was extracted (mean = 54%) in the first two steps of the SEP (Figure 7.2; Table 7.3).

Arsenic recovered upon the dissolution of crystalline Fe oxides in the Step-IV constituted $\leq 24\%$ of the total soil As, except for the Faulkners, Dunoon, Wollongbar and Adelaide Hills samples where it was $\geq 34\%$. The crystalline Fe content in these four samples was $\geq 87.5\%$ of the free Fe content (see Table 7.1), therefore it could be expected that As was adsorbed or coprecipitated in greater proportion with the crystalline Fe oxides in these soils (Table 7.3). The residual phase of As was $\leq 11.7\%$ of the total As in the cattle-dip site and railway corridor soils, while no residual As was found in the As spiked soils (Figure 7.2). Low amount of residual As in historically contaminated soils reflect that these soils could potentially release As in the environment due to a greater proportion of As being present in the potentially mobilisable phases.

The differences in As partitioning between the As spiked and the long term contaminated (cattle-dip sites and railway corridor) soils (Figure 7.2, Table 7.3) possibly occurred due to the variation in the equilibration or ageing time of the soils (Tang et al. 2007; Smith et al. 2008). In spiked soils a much higher proportion of As was removed in the Step-I and Step-II (mean = 12.7 and 41%) of the SEP than that of the long term contaminated soils (mean = 1.2 and 13%). This could be attributed to the shorter equilibration time of added As as As^{V} in the spiked soils compared to the prolonged ageing (over 40 years) in the contaminated soils (Smith et al. 2008).

Amorphous Fe ($r = 0.63$; $P < 0.001$) and organic carbon ($r = 0.40$; $P < 0.05$) in the contaminated and spiked soils were positively correlated with amorphous Fe oxides bound As (Step-III). The disparities in As speciation among the four artificially As-contaminated soils were related to their varying soil properties, such as pH, free Fe and clay contents. For

example, lower As in the exchangeable phase of the Westwood soil ($\leq 2.2\%$) was attributed to the higher free Fe content (3.3 %) in this sample. Conversely, higher As removal in the Step-II of Narrabri soil could be due to its relatively lower free Fe content (1.28 %), alkaline pH (7.53) and higher clay content (60%) (Smith et al. 1998; Goldberg 2002). Similarly, Juhasz et al. (2008) reported that higher Fe content and lower pH in a Red Ferrosol (9.96 % Fe, pH 5.8) than that of a Brown Chromosol (0.8 % Fe, pH 7.8) reduced As bioavailability up to 75%, after 12-month of incubation. The results from this study also indicate that soil properties mainly free Fe content and pH appeared to affect the distribution and phytoavailability of As in the spiked soils, albeit all spiked soils were equilibrated for same time (Wenzel et al. 2001; Novoa-Munoz et al. 2007).

7.3.3 Arsenic speciation using XANES spectroscopy

The As K-edge XANES spectra of the reference compounds and selected soil samples are presented in Figure 7.3a,b. The white line energy position of As (As^{V} and As^{III}) sorbed on Fe oxides and As present in mineral forms was in the order: scorodite > $\text{As}^{\text{V}}\text{-Fh}$, $\text{As}^{\text{V}}\text{-Cop-Fh}$ > $\text{As}^{\text{V}}\text{-Gt}$ > $\text{As}^{\text{III}}\text{-Fh}$ > As_2S_3 (Figure 7.3a). The LCF results from XANES spectra of soil samples showed that As was predominantly present in As^{V} form and mainly associated with the amorphous Fe oxides (14–78% of the total As) (Table 7.4). Arsenic was depicted to present as $\text{As}^{\text{V}}\text{-Fe}^{\text{III}}$ coprecipitates in two samples, Piora (64%) and Murray Bridge (28%). Previous studies on As speciation in the As-contaminated mining and pesticides manufacturing sites also showed the predominance of As^{V} associated with amorphous Fe oxides (Cances et al. 2005; Arai et al. 2006; Meunier et al. 2010). Arsenic in this fraction is considered to be a potential source of As release to the environment as indicated by Meunier et al. (2010), who investigated As speciation in tailings and mining-impacted soils.




Figure 7.3 Arsenic K-edge XANES spectra of (a) reference compounds, and (b) selected soil samples from the cattle-dip sites, railway corridor and As spiked soils. Solid (—), dotted (···) and dashed (---) lines drawn at 11875.7, 11872.2 and 11868.7 eV represent the position of white lines for arsenate (As^{V}), arsenite (As^{III}), and orpiment ($\text{As}_2^{\text{III}}\text{S}_3$), respectively. The white line energy values (eV) are given in the parentheses for each of the reference compound.

They measured bioaccessibility of As in gastric and intestinal phases and reported higher As bioaccessibility (~10%) in the samples having greater proportion of As-bearing amorphous phases (amorphous Fe oxides or calcium-Fe-arsenate) than those where As was present in arsenopyrite and scorodite (< 1%) minerals. The XANES and SEP (Step-III) data also showed the greatest proportion (51% and 75% of total As) of As being present in the potentially available (As-bearing amorphous Fe oxides) form (Figure 7.2; Table 7.4). In the current study, higher proportion of As was found in this phase than those reported by Meunier et al. (2010) (i.e. 2–7%) during dissolution of As-bearing amorphous forms, which could be attributed to the differences in source of As contamination in the two studies, as mentioned earlier.

LCF results showed that > 95% of the total As was present in association with the amorphous Fe oxides in the As spiked soils (Figure 7.3; Table 7.4); the Lansdowne samples were excluded from the analysis because of their low As contents. These results are different to the SEP data where upon dissolution of the amorphous Fe oxides (Step-III) 22–57% (mean = 41.7%) of the total As was extracted from the three spiked soils.

Table 7.4 Linear least-squares combination fits of XANES spectra of reference compounds with those of soil samples ($n = 30$) analyzed. Fit results correspond to XANES spectra of selected soil samples plotted in Figure 7.3.

Soil samples*	Reference compounds (% weights) [†]				R-factor ^e
	As ^V -Fh ^a	As ^V -Gt ^b	As ^{III} -Fh ^c	Scorodite ^d	
<i>Cattle-dip sites (n = 18)</i>					
Dungarubba	57	4	35	4	0.0034
Hendersons	61	6.5	30	2.6	0.0022
Piora	26 (64) ^f	–	10	–	0.0033
Sneesbyslane	31	69	–	–	0.0040
JBM	92	–	8	–	0.0041
Boorie	34	66	–	–	0.0019
Henrywhite	48	14	32	6	0.0031
Enak	96	–	–	4	0.0005
Faulkners	67		20	13	0.0041
Dunoon	79	21	–	–	0.0033
Nimbin	78	–	16	6	0.0028
Blackwoods	81	19	–	–	0.0049
Blackwoods30–40 cm	89	11	–	–	0.0023
Alexanders	96	–	–	4	0.0005
Goolmangar	80	–	–	20	0.0016
Sandyarm	61 (17) ^g	–	–	22	0.0066
Sandyarm30–40 cm	14 (59) ^g	3	24	–	0.0083
Wollongbar	81	–	–	19	0.0008
<i>Railway corridor (n = 3)</i>					
Murray bridge	52 (28) ^f	–	20	–	0.0035
Adelaide hills	96	4	–	–	0.0008
Kapunda	56	6	38	–	0.0041
<i>As spiked soils (n = 9)</i>					
Westwood-100	98	–	–	–	0.0006
Westwood-200	97	–	–	–	0.0006
Westwood-400	100	–	–	–	0.0005

Table 7.4 Continued...

Soil samples*	Reference compounds (% weights) [†]				R-factor ^e
	As ^V -Fh ^a	As ^V -Gt ^b	As ^{III} -Fh ^c	Scorodite ^d	
Narrabri-100	96	–	4	–	0.0177
Narrabri-200	98	–	2	–	0.0007
Narrabri-400	95	–	–	–	0.0032
Lucerne-100	100	–	–	–	0.0003
Lucerne-200	96	4	–	–	0.0005
Lucerne-400	97	3	–	–	0.0014

^aAs^V adsorbed to ferrihydrite; ^bAs^V adsorbed to goethite; ^cAs^{III} adsorbed to ferrihydrite;

^dMineral phase of As^V-Fe^{III} (FeAsO₄·2H₂O); ^fAs^V coprecipitated ferrihydrite (As^V-Cop-Fh);

^gOrpiment (As₂S₃); ^eNormalized sum of the residuals (R-factor) = $\sum ((y - y_{fit})^2) / \sum (y^2)$.

*All soil samples are collected at 0–10 cm depth, except for the two samples mentioned at 30–40 cm.

[†]LCF results are presented with < ±5% delta weights

The results indicate that XANES spectroscopy may overlook the exchangeable and specifically sorbed fractions of As in historically contaminated and spiked soils. While the SEP allowed to determine these two highly mobile As phases in the studied soils, notably in the As spiked soils where more than 50% As was extracted in the first two fractions of the SEP. XANES spectra of two soils (Boorie and Wollongbar) was also analysed after Step-II and Step-III of the SEP to elucidate any change in the spectral features after removal of As in respective steps of the SEP (Figure 7.4). The data for these two soils (after Step-II and Step-III) show nearly identical XANES spectra.




Figure 7.4 Arsenic K-edge XANES spectra for Boorie (black and purple lines) and Wollongbar (red and green lines) soil samples analysed after Step-II and Step-III of the SEP. The solid (–) line is drawn at 11875.7 eV represent the position of white line for arsenate (As^{V}).

The lack of a reference compound representing only the exchangeable fraction of As might be one of the drawbacks in the XANES analysis. The results suggest that combined use of SEP and XANES is a powerful approach to determine the presence of more labile (and toxic) forms of As in soils which could be underestimated using XANES spectroscopy alone.

Arsenate was also detected to present in adsorbed form onto the surface of crystalline Fe oxides (mean = 19%) (Table 7.4). In addition to the presence of As^{V} -Fe oxides adsorption

complexes, a mineral phase of As^{V} , scorodite ($\text{FeAsO}_4 \cdot 2\text{H}_2\text{O}$) was also identified in 10 of the 18 samples from cattle-dip sites. Scorodite has been previously reported to occur mostly in the As-contaminated mining sites (Paktunc et al. 2003; Meunier et al. 2010).

Arsenic occurred in As^{III} oxidation state in association with the amorphous Fe oxides in the cattle-dip sites and railway corridor samples (Table 7.4), with the maximum value of 38% observed in Kapunda soil (Figure 7.3). Arsenite was detected in two of the As spiked soils, where it was < 5% of the total As. Reduction of As^{V} to As^{III} due to the high intensity synchrotron X-rays was discounted considering the low levels of As^{III} observed in the As spiked soils. In a previous study, Smith et al. (2006b) used a chemical method for As speciation in herbicide impacted railway corridor soils and reported up to 40% of the total As in As^{III} form. They postulated that regular application of As^{III} -based herbicides may have led to the increased sorption of As^{III} over As^{V} on the surface of amorphous Fe oxides under alkaline ($\text{pH} > 7$) soil conditions. This mechanism may explain the presence of As^{III} in the two railway corridor As^{III} -herbicides contaminated soil samples (see Table 7.4) (Goldberg 2002; Smith et al. 2006b). The presence of As^{III} in the oxidised and acidic soil conditions in the As^{III} -pesticides impacted cattle-dip site soils (see Table 7.4) could possibly be attributed to the microbial reduction of As^{V} to As^{III} as reported by Turpeinen et al. (2002). These authors suggested that microbes could transform As^{V} to As^{III} under aerobic conditions in As-contaminated soils with acidic pH (4.4–4.7). Reduction of As^{V} can also occur due to complex redox environments that exist in rhizosphere soils. However, it is not immediately evident what factors contributed toward the occurrence of As in As^{III} form only in these soils; further site-specific detailed examinations are warranted to elucidate the factors controlling the oxidation-reduction of As in soils.

A decrease in the white line energy position (11870.03 eV) in the Sandyarm sample collected from 30–40 cm depth (see Figure 7.3b) indicated the presence of orpiment-like precipitates (As_2S_3 , 59%), an As^{III} -sulfide mineral phase of As. The presence of As in this form might be due to the presence of sulfur added through superphosphate fertilisers. The presence of microbes and prevailing reducing conditions in the sub-surface soil may have led to the formation of $\text{As}^{\text{III}}-\text{S}^{2-}$ species as reported by Arai et al. (2006) for a contaminated site near to an arsenical pesticides manufacturing plant.

7.3.4 Arsenic phytoavailability to *B. juncea* in relation to SEP and XANES spectroscopy

Arsenic concentration in the shoot of *B. juncea* ranged from 3.6–74.8 mg kg^{-1} with the maximum value (74.8 mg kg^{-1}) observed in Blackwoods30–40 cm sample (Table 7.3). The shoot As concentration varied among the studied samples depending upon the degree of contamination and plant availability of As in the soils. The lighter textured (clay $\leq 6\%$) soils with low free Fe content ($\leq 0.22\%$) including Dungarubba, Sandyarm, Sandyarm30–40 cm exhibited relatively greater plant As concentration, although the total As concentration in these soils was much lower than the many other contaminated soils (Table 7.3).




Figure 7.5 Arsenic concentration in dry shoots of *B. juncea* in relation to the sequentially extracted (a) exchangeable As (Step-I); (b) specifically sorbed As (Step-II); (c) amorphous Fe oxides bound As (Step-III); (d) crystalline Fe oxides bound As (Step-IV); and (e) As determined using XANES associated with amorphous Fe oxides. Cattle-dip sites and railway corridor soils (●); As spiked soils (●). The correlations are presented excluding the As spiked soils data due the considerable differences in ageing time.

The concentration of As in the plant shoot was found to be greater in all samples than the normally reported value of $1.5 \mu\text{g kg}^{-1}$ for plants grown in uncontaminated soils (McLaren et al. 2006). The phytotoxicity symptoms of As such as marked reduction in plant growth were also observed across all the contaminated and As spiked soils (Sheppard 1992; Jiang and Singh 1994). Similarly, Jiang and Singh (1994) reported a substantial decrease in the dry biomass yield of rye grass and barley plants due to the application of As, which was attributed to the inhibitory effect of As (as As^{V}) on the phosphate uptake pathway leading to insufficient levels of phosphorylated compounds. Arsenic in As^{III} form is known to have twice as much phytotoxicity as As^{V} , since it rapidly combines with the dithiol functional groups and destroys the functioning of sulfhydryl enzymes; thereby causing the membrane degradation and immediate cell death. There was greater As toxicity that severely affected plant growth (qualitative observations) and markedly reduced dry matter yield (Appendix 4) of *B. juncea* plants in Dungarubba, Sandyarm and Sandyarm30–40 cm soils (Sheppard 1992; Mandal and Suzuki 2002). This supports the findings that As in these particular soils exists in highly toxic (As^{III}) and plant available form (Table 7.4).

The shoot As concentration of *B. juncea* in all As spiked samples was comparatively greater than the long term contaminated soil samples which agrees with the SEP data showing greater proportion of As in the first two steps (Table 7.3). Among the As spiked samples, maximum shoot As concentration of 44.5 mg kg^{-1} was observed in Lansdowne-20 followed by Lucerne-200 and Narrabri-200 samples (Table 7.3). The plants in Narrabri-400 and Lucerne-400 soils died immediately after germination as a result of severe As toxicity at these high As levels (Sheppard 1992). The lower shoot As concentration observed in the Westwood samples

(6.4–12.4 mg kg⁻¹) compared to the other three As spiked samples could be related to the minimum amount of As in the exchangeable phase in this soil.




Figure 7.6 The relationship of amorphous (ammonium oxalate extractable) Fe concentration with the amorphous Fe oxides associated As in the studied soils determined using (a) SEP in Step-III, and (b) XANES spectroscopy.

The concentration of As in the plant shoot showed a highly significant ($P < 0.001$) positive correlation with the exchangeable As ($r = 0.85$), indicating that As in this phase is readily available to plants (Figure 7.5a). The results are in agreement with those of Anawar et al. (2008), who also reported a significant positive relationship ($r = 0.94$; $P < 0.05$) between

exchangeable As and plant As concentration in alfalfa. Arsenic concentration in shoots was also significantly correlated with the amorphous Fe oxides bound As fraction estimated by SEP in Step-III ($r = 0.67$; $P < 0.01$) and XANES ($r = 0.54$; $P < 0.05$) (Figure 7.5c,e). The experimental data indicate that As associated with amorphous Fe oxides was also plant available. A significant positive correlation was observed between the amorphous Fe oxide content and As associated with amorphous Fe oxides fraction estimated using SEP (Step-III, $r = 0.63$; $P < 0.01$) and XANES ($r = 0.50$; $P < 0.01$) (Figure 7.6), which further supports the earlier suggestion. The phytoavailability of As from this fraction could possibly be attributed to the release of As from amorphous Fe oxides phase, since this As fraction is considered less stable than those bound with the crystalline forms (Juhasz et al. 2007; Juhasz et al. 2009). Arsenic can be released or solubilised from this fraction due to the effects of low-molecular-weight citric and oxalic acids excreted by plant roots, and/or changes in soil pH and redox conditions (Smith et al. 1998; Tao et al. 2006; Slowey et al. 2007; Mikutta et al. 2010).

7.4 CONCLUSIONS

The results show that combination of SEP and XANES techniques provided vital information on the speciation of As in soils, which could be useful to better understand the lability, toxicity and phytoavailability of As in soils. XANES spectroscopy identified oxidation state ($\text{As}^{\text{V}}/\text{As}^{\text{III}}$) and mineral phases (e.g. scorodite, orpiment and coprecipitated or adsorbed forms) of As in soils, whereas the SEP was better in quantifying the highly mobile (Step-I and Step-II) As pools in soils.

Arsenic existed in soluble and mobilisable forms, and both As^{V} and As^{III} forms in historically contaminated soils; and the plant As uptake data reflect that As in these forms is

phytoavailable. Soil properties especially Fe oxides and organic carbon contents are important in governing the speciation and phytoavailability of As in contaminated soils. Such detailed knowledge is vital for the reclamation and management of these As-contaminated soils.

7.5 REFERENCES

- Anawar HM, Garcia-Sanchez A, Regina IS (2008) Evaluation of various chemical extraction methods to estimate plant-available arsenic in mine soils. *Chemosphere* 70 (8):1459-1467.
- Arai Y, Lanzirrotti A, Sutton SR, Newville M, Dyer J, Sparks DL (2006) Spatial and temporal variability of arsenic solid-state speciation in historically lead arsenate contaminated soils. *Environ Sci Technol* 40 (3):673-679.
- Cances B, Juillot F, Morin G, Laperche V, Alvarez L, Proux O, Hazemann JL, Brown GE, Calas G (2005) XAS evidence of As(V) association with iron oxyhydroxides in a contaminated soil at a former arsenical pesticide processing plant. *Environ Sci Technol* 39 (24):9398-9405.
- Gillman GP, Bruce RC, Davey BG, Kimble JM, Searle PL, Skjemstad JO (1983) A comparison of the methods used for determination of cation exchange capacity. *Commun Soil Sci Plant Anal* 14 (11):1005-1014.
- Goldberg S (2002) Competitive adsorption of arsenate and arsenite on oxides and clay minerals. *Soil Sci Soc Am J* 66 (2):413-421.
- Grafe M, Nachtegaal M, Sparks DL (2004) Formation of metal-arsenate precipitates at the goethite-water interface. *Environ Sci Technol* 38 (24):6561-6570.
- Huang PM, Fujii R (1996) Selenium and arsenic. In: Sparks DL (ed) *Methods of Soil Analysis. Part 3, Chemical Methods*. Soil Science Society of America Inc, Wisconsin, pp 793-831.
- Jiang QQ, Singh BR (1994) Effect of different forms and sources of arsenic in crop yield and arsenic concentration. *Water Air Soil Pollut* 74 (3-4):321-343.

- Juhasz AL, Smith E, Weber J, Naidu R, Rees M, Rofe A, Kuchel T, Sansom L (2008) Effect of soil ageing on in vivo arsenic bioavailability in two dissimilar soils. *Chemosphere* 71 (11):2180-2186.
- Juhasz AL, Smith E, Weber J, Rees M, Rofe A, Kuchel T, Sansom L, Naidu R (2007) In vitro assessment of arsenic bioaccessibility in contaminated (anthropogenic and geogenic) soils. *Chemosphere* 69 (1):69-78.
- Juhasz AL, Weber J, Smith E, Naidu R, Rees M, Rofe A, Kuchel T, Sansom L (2009) Assessment of four commonly employed in vitro arsenic bioaccessibility assays for predicting in vivo relative arsenic bioavailability in Contaminated Soils. *Environ Sci Technol* 43 (24):9487-9494.
- Kachenko AG, Grafe M, Singh B, Heald SM (2010) Arsenic speciation in tissues of the hyperaccumulator *P. calomelanos* var. *austroamericana* using X-ray absorption spectroscopy. *Environ Sci Technol* 44 (12):4735-4740.
- Kimber SWL, Sizemore DJ, Slavich PEG (2002) Is there evidence of arsenic movement at cattle tick dip sites? *Aust J Soil Res* 40 (7):1103-1114.
- La Force MJ, Fendorf S (2000) Solid-phase iron characterization during common selective sequential extractions. *Soil Sci Soc Am J* 64 (5):1608-1615.
- Lombi E, Sletten RS, Wenzel WW (2000) Sequentially extracted arsenic from different size fractions of contaminated soils. *Water Air Soil Pollut* 124 (3-4):319-332.
- Mandal BK, Suzuki KT (2002) Arsenic round the world: a review. *Talanta* 58 (1):201-235.
- Manning B (2005) Arsenic speciation in As(III)- and As(V)-treated soil using XANES spectroscopy. *Microchim Acta* 151 (3-4):181-188.

- McLaren RG, Megharaj M, Naidu R (2006) Fate of arsenic in the soil environment. In: Naidu R, Smith E, Owens G, Bhattacharya P, Nadebaum P (eds) *Managing Arsenic in the Environment: From Soil to Human Health*. CSIRO, Collingwood, Victoria, pp 157-182.
- McLaren RG, Naidu R, Smith J, Tiller KG (1998) Fractionation and distribution of arsenic in soils contaminated by cattle dip. *J Environ Qual* 27 (2):348-354.
- Mehra OP, Jackson ML (1960) Iron oxide removal from solid clays by a dithionite-citrate system buffered with sodium bicarbonate. *Clays Clay Miner* 7:317-327.
- Meunier L, Walker SR, Wragg J, Parsons MB, Koch I, Jamieson HE, Reimer KJ (2010) Effects of soil composition and mineralogy on the bioaccessibility of arsenic from tailings and soil in gold mine districts of Nova Scotia. *Environ Sci Technol* 44 (7):2667-2674.
- Mikutta C, Frommer J, Voegelin A, Kaegi R, Kretzschmar R (2010) Effect of citrate on the local Fe coordination in ferrihydrite, arsenate binding, and ternary arsenate complex formation. *Geochim Cosmochim Acta* 74 (19):5574-5592.
- Miller RO (1998) Nitric-perchloric acid wet digestion in an open vessel. *Handbook of Reference Methods for Plant Analysis*, Y.P. Kalra edn. CRC Press, Boca Raton, Florida, U.S.A.
- National Environment Protection (Assessment of Soil Contamination) Measure (NEPM) (1999) *Investigation Levels for Soil and Groundwater*. National Environmental Protection Council Service Corporation, Adelaide, Australia.
- Novoa-Munoz JC, Queijeiro JMG, Blanco-Ward D, Alvarez-Olleros C, Garcia-Rodeja E, Martinez-Cortizas A (2007) Arsenic fractionation in agricultural acid soils from NW Spain using a sequential extraction procedure. *Sci Total Environ* 378 (1-2):18-22.

- Ostergren JD, Brown GE, Parks GA, Tingle TN (1999) Quantitative speciation of lead in selected mine tailings from Leadville, CO. *Environ Sci Technol* 33 (10):1627-1636.
- Paktunc D, Foster A, Laflamme G (2003) Speciation and characterization of arsenic in Ketz River mine tailings using x-ray absorption spectroscopy. *Environ Sci Technol* 37 (10):2067-2074.
- Payne RW, Murray DA, Harding SA, Baird DB, Soutar DM (2009) GENSTAT for Windows - Introduction. 12th edn edn. VSN International, Hemel Hempstead, UK.
- Ravel B, Newville M (2005) ATHENA, ARTEMIS, HEPHAESTUS: data analysis for X-ray absorption spectroscopy using IFEFFIT. *J Synchrotron Rad* 12:537-541.
- Rayment GE, Lyons DJ (2011) Soil Chemical Methods - Australasia. CSIRO Publishing, Collingwood, VIC, Australia.
- Scheinost AC, Kretzschmar R, Pfister S, Roberts DR (2002) Combining selective sequential extractions, X-ray absorption spectroscopy, and Principal Component analysis for quantitative zinc speciation in soil. *Environ Sci Technol* 36 (23):5021-5028.
- Shah P, Strezov V, Stevanov C, Nelson PF (2007) Speciation of arsenic and selenium in coal combustion products. *Energy Fuels* 21 (2):506-512.
- Sheppard SC (1992) Summary of phytotoxic levels of soil arsenic. *Water Air Soil Pollut* 64:539-550.
- Slowey AJ, Johnson SB, Newville M, Brown GE (2007) Speciation and colloid transport of arsenic from mine tailings. *Appl Geochem* 22 (9):1884-1898.
- Smith E, Naidu R, Alston AM (1998) Arsenic in the soil environment: A review. *Adv Agron* 64:149-195.

- Smith E, Naidu R, Weber J, Juhasz AL (2008) The impact of sequestration on the bioaccessibility of arsenic in long-term contaminated soils. *Chemosphere* 71 (4):773-780.
- Smith E, Owens G, Naidu R (2006a) Arsenic in the Australian environment. In: Naidu R, Smith E, Owens G, Bhattacharya P, Nadebaum P (eds) *Managing Arsenic in the Environment. From Soil to Human Health*. CSIRO, Collingwood, Victoria, pp 627-638.
- Smith E, Smith J, Naidu R (2006b) Distribution and nature of arsenic along former railway corridors of South Australia. *Sci Total Environ* 363:175-182.
- Tang XY, Zhu YG, Shan XQ, McLaren R, Duan J (2007) The ageing effect on the bioaccessibility and fractionation of arsenic in soils from China. *Chemosphere* 66 (7):1183-1190.
- Tao YQ, Zhang SZ, Jian W, Yuan CG, Shan XQ (2006) Effects of oxalate and phosphate on the release of arsenic from contaminated soils and arsenic accumulation in wheat. *Chemosphere* 65 (8):1281-1287.
- Turpeinen R, Panssar-Kallio M, Kairesalo T (2002) Role of microbes in controlling the speciation of arsenic and production of arsines in contaminated soils. *Sci Total Environ* 285 (1-3):133-145.
- Voegelin A, Weber FA, Kretzschmar R (2007) Distribution and speciation of arsenic around roots in a contaminated riparian floodplain soil: Micro-XRF element mapping and EXAFS spectroscopy. *Geochim Cosmochim Acta* 71 (23):5804-5820.
- Wenzel WW, Kirchbaumer N, Prohaska T, Stingeder G, Lombi E, Adriano DC (2001) Arsenic fractionation in soils using an improved sequential extraction procedure. *Anal Chim Acta* 436 (2):309-323.