

1 **Morphological changes of the lateral meniscus in end**
2 **stage lateral compartment osteoarthritis of the knee**

3
4 Seung Hyun Hwang, MD[†], Kwang Am Jung MD[†], Won Jun Lee MD[†], Ki Hyuk

5 Yang MD[‡], Dong Won Lee, RS^{†§}, Aaron Carter MD^{||},

6 David John Hunter MBBS PhD^{¶††}

7
8 [†] Joint & Arthritis Research, Department of Orthopaedic Surgery, Himchan
9 Hospital, Seoul, Korea

10 [‡] Department of Diagnostic Radiology, Himchan Hospital, Seoul, Korea

11 [§] The Webb school of California, USA

12 ^{||} Rothman Institute, Philadelphia PA, USA

13 [¶] Division of Research, New England Baptist Hospital, Boston MA, USA

14 ^{††} Rheumatology Department, Royal North Shore Hospital and Northern Clinical
15 School, University of Sydney, NSW Australia.

16 **Keywords:** Lateral meniscus, osteoarthritis

17
18 **Reprint requests to** Kwang Am Jung, MD

19 **Address:** Joint and Arthritis Research, Department of Orthopaedic Surgery, Himchan Hospital,
20 20-8, Songpa-dong, Songpa-gu, Seoul, Korea, Email: kwangamj@gmail.com

21 **Abstract**

22 **Objective:** The aim of this study was to evaluate the morphological changes of
23 the lateral meniscus in end stage lateral compartment osteoarthritis(OA) of the
24 knee.

25 **Methods:** 158 knee joints from 133 patients that subsequently underwent total
26 knee joint arthroplasty from January 2008 to December 2009 were enrolled.
27 There were 26 men and 107 women. Their ages ranged from 56 to 81 (mean
28 67.4 ± 6.5 yrs). All study participants had complete obliteration of the lateral joint
29 space identified by weight bearing radiography. Meniscal position was assessed
30 by measuring meniscal subluxation and meniscal height. The meniscal
31 morphology was assessed using a modified whole-organ magnetic resonance
32 imaging score (WORMS). The frequency of different meniscal morphologies
33 and their respective positions were calculated.

34 **Results:** The predominant type (42.4, 53.8 and 52.5% in the anterior horn, mid
35 body and posterior horn, respectively) of abnormal meniscal morphology was a
36 complete maceration/destruction or complete resection. The anterior horn of
37 non-macerated lateral meniscus was more subluxed than that of the non-
38 macerated medial meniscus in patients with lateral osteoarthritis.

39 **Conclusion:** This study suggests that the lateral meniscus in persons with end
40 stage lateral osteoarthritis are mostly macerated or destroyed. Also, unlike
41 isolated end staged medial compartment osteoarthritis, the anterior horn of the
42 lateral meniscus in isolated end stage lateral osteoarthritis is commonly affected.

43 **Key words:** Meniscus, Lateral Osteoarthritis

44

45 **Introduction**

46 Structural changes in knee osteoarthritis are characterized by significant
47 cartilage loss, subchondral sclerosis, osteophytosis, subchondral cysts,
48 meniscal degeneration, and other intraarticular or extraarticular soft tissue
49 abnormalities[1-5]. In addition to extensive investigation of the biology and
50 genetic etiology of osteoarthritis [6-9], investigators have attempted to describe
51 the morphological characteristics associated with the above structural
52 changes[10-14]. Among them, the meniscus, as one of the soft tissues
53 prominently involved in OA etiopathogenesis, has been evaluated based on its
54 integral role in knee function[15-17]. Several studies have shown that both
55 meniscal subluxation and meniscal tears are common not only in knee OA, but
56 particularly frequent in knees with radiographic knee OA and appear to be
57 related to the degree of joint space narrowing on plain radiographs[11, 15-16,
58 18-19] Based on prior reports and existing dogma[11], the common consensus
59 is that advanced stage OA of the knee, with complete loss of either the medial
60 or lateral compartment joint space on radiographs, might be associated with a
61 completely macerated/destroyed meniscus and hyaline cartilage. However, a

62 previous published study found no correlation between the radiological and
63 morphological changes of the medial meniscus in end stage medial
64 osteoarthritis[20], where a hypertrophied meniscus was the most prevalent
65 finding. Another recent study showed that OA knees have thicker menisci than
66 those of non-OA knees[21] . In terms of lateral tibiofemoral arthritis, to date,
67 there is one study comparing the prevalence of lateral tibiofemoral (TF)
68 osteoarthritis in Asian and western population[22] suggesting that Asian knees
69 have more lateral TF OA. What accounted for this lateral TF increase in Asian
70 knees is not currently known. Furthermore there is little known of the
71 morphological and positional changes of the lateral meniscus in patients with
72 advanced lateral OA. Therefore, the aim of this study was to examine the
73 morphological and positional changes of the lateral meniscus in patients with
74 advanced lateral compartment OA.

75

76 **Materials and Methods**

77

78 **Study participants**

79 The research design used in this study was a consecutive series of
80 persons presenting with end-stage lateral tibiofemoral (TF) OA to an orthopedic
81 specialty hospital. 143 potential patients participated in this study. All potential

82 patients presented with lateral compartment osteoarthritis. A series of knee
83 radiographs (weight bearing posteroanterior radiographs, weight bearing 30
84 degrees posteroanterior, lateral, and skyline views) were obtained for each
85 patient to determine whether advanced lateral TF compartment radiographic OA
86 was present. The radiographs were graded using the Kellgren-Lawrence (K&L)
87 grading scale[23] and scored for lateral joint space narrowing (JSN) on a scale
88 of 0 (normal)—3 (total loss of the joint space) with the help of the Osteoarthritis
89 Research Society International (OARSI) atlas[24] by two experienced clinicians
90 (SHH, WJL with 7 and 6 years of musculoskeletal radiology experience respectively)..
91 If patients had complete lateral joint space obliteration (K&L = 4 and JSN = 3)
92 on the weight-bearing posteroanterior radiograph, they were eligible for this
93 study. The inter-rater reliability of **KL and OARSI grading** were determined by
94 calculating the intraclass correlation coefficients (ICCs), which was 0.96 and
95 0.95, respectively. 10 patients were excluded due to the diagnosis of secondary
96 OA (i.e., OA associated with fracture, prior knee associated arthroscopic or
97 open surgery, or another disease process), simultaneous medial compartment
98 OA with medial joint space narrowing, and systemic inflammatory arthritis.(e.g.,
99 rheumatoid arthritis, gout), based on medical records. Patients without
100 contraindications to MRI underwent MRI of their abnormal knee joints. Finally, a
101 total of 158 knee joints among 133 patients were included for this study from
102 January, 2008 to December 2009. The study was approved by the hospital
103 ethics committee at our institution and all patients gave written informed
104 consent to use their anonymised data

105

106 **Limb alignment assessment**

107 The degree of valgus deformity was measured as the femorotibial angle (FTA)
108 by two experienced raters (SHH, WJL) using a standing long limb radiograph.
109 Femorotibial angles (FTA) were measured by drawing a line along the axis of
110 the femoral shaft to intersect the corresponding line drawn through the tibial
111 shaft. During the assessment, the readers were blinded to MRI results. The
112 inter-rater reliability of the FTA measurements were determined by calculating
113 the intraclass correlation coefficients (ICCs), which was 0.93.

114

115 **MRI acquisition** Meniscus changes were assessed using a 1.5 T MRI system in
116 the sagittal and coronal planes with spin-echo (proton density weighted
117 acquisition) and fast scan (T2-weighted images) techniques. The sagittal and
118 coronal spin-echo proton density weighted acquisition images were acquired
119 using the following parameters: 1800/15/2(TR/TE/NEX), slice thickness 4 mm,
120 inter-slice gap 0.4 mm for coronal images and 0.3 mm for sagittal images, slice
121 thickness 3 mm, and matrix 256 X 256. T2-weighted images were also acquired
122 using the following parameters: 3700/100/2(TR/TE/NEX), slice thickness 4 mm,
123 inter-slice gap 0.4 mm for coronal images, and a slice thickness of 3 mm with
124 an inter-slice gap of 0.3 mm for sagittal images.

125

126 **MRI interpretation**

127 The anterior and posterior horns and mid-bodies of menisci were examined for
128 (1) meniscal morphology, and (2) meniscal position (Figs. 1, 2). During the
129 assessment, the readers were blinded to radiographic results, patient symptoms,
130 patient age, and other clinical data. Meniscal morphology (integrity) was
131 measured independently by two experienced observers (SHH, WJL), and the
132 overall ratings were determined by consensus when necessary. The
133 morphology at each portion of the meniscus was assessed, using a modification
134 of the whole-organ magnetic resonance imaging score (WORMS) assessment
135 system [25]. According to the modifications reported in a previous study[20]: 0 =
136 intact, 1 = minor radial tear or parrot-beak tear, 2 = non-displaced tear, 3 =
137 displaced but no tear, 4 = displaced tear or partial resection, 5 = hypertrophied
138 displaced, 6 = hypertrophied displaced tear, and 7 = complete
139 maceration/destruction or complete resection (Fig. 1). The meniscal integrity of
140 the anterior and posterior horns of the menisci was measured in the sagittal and
141 coronal planes, in which the meniscal morphology was best observed. The mid-
142 body height was measured where the medial and lateral tibial spine volume was
143 maximal [11-12, 17]. “Hypertrophy” was considered present when the lateral
144 meniscus height was 2 mm greater than the medial meniscus, regardless of the
145 medial meniscus width, using reference values of the normal meniscus height in
146 which those of the lateral meniscus are normally smaller than the medial
147 meniscus[17] . The inter-rater reliability of meniscal morphology ratings was
148 0.87 (kappa) for meniscal morphology at the anterior horn of the lateral
149 meniscus, 0.80 at the mid-body of the lateral meniscus, and 0.86 at the
150 posterior horn of the lateral meniscus.

151 Two experienced observers (SHH, WJL) independently measured the
152 meniscal position; the mean values were used for the analysis. The meniscal
153 position was assessed by measuring the meniscal subluxation and height of
154 each knee (Figure 2). To determine the meniscal height, the anterior and
155 posterior horns of the menisci were measured in the sagittal plane, which
156 allowed the best visualization of the greatest meniscal size. The mid-body
157 height was measured in the coronal plane, where the medial and lateral tibial
158 spine volume was maximal. The meniscal height was measured at the most
159 peripheral edge of each meniscus, regardless of whether the meniscus was “in-
160 place”, subluxed or extruded. To determine meniscal subluxation, anterior
161 subluxation of the anterior horn of the medial and lateral meniscus was
162 assessed in the area where the subluxation was most prominent through
163 multiple sagittal slices. Lateral subluxation of the mid-body of the lateral and
164 medial subluxation of the medial meniscus was measured where the volume of
165 the medial and lateral tibial spine was greatest. Posterior subluxation of the
166 posterior horn was not measured, because this could not be performed
167 accurately in the sagittal plane. For completely macerated or destroyed menisci,
168 meniscal subluxation and the meniscal height could not be measured (Fig. 2)
169 and were handled as missing values and 0 mm, respectively, for statistical
170 analysis. The inter-rater reliability of the meniscal position measurements were
171 determined by calculating the intraclass correlation coefficients (ICCs). An ICC
172 of 1 suggests perfect reliability, and an ICC > 0.75 and ICC < 0.4 is generally
173 considered to represent excellent and poor reliability respectively. For the cases
174 that could be measured, the ICCs for the meniscal height and meniscal

175 subluxation were the height at the anterior horn 0.74, mid-body 0.81, and
176 posterior horn 0.80; anterior subluxation at the anterior horn 0.85, and lateral
177 subluxation at the mid-body 0.84.

178

179 **Statistical analysis**

180 First, the frequency of the meniscal morphology for each portion of the
181 meniscus was determined. Second, the medial meniscus height and
182 subluxation in cases with a non-macerated lateral meniscus were compared
183 with those in cases with a macerated lateral meniscus using the independent T-
184 test. Third, the lateral meniscus height and subluxation in cases with a non-
185 macerated lateral meniscus were compared to those of the medial meniscus
186 using the paired sample t test at each meniscal portion, including the anterior
187 horn, midbody, and posterior horn. The differences in FTA valgus angle was
188 evaluated according to each meniscal morphological type at anterior, midbody
189 and posterior horn, respectively with one-way analysis of variance with post hoc
190 comparison (Turkey's test). For height and subluxation in the cases with a
191 macerated lateral meniscus, where each value was unmeasurable, comparison
192 with the cases that had a non-macerated meniscus could not be performed.
193 Correlations between the BMI (Body mass index), FTA (femorotibial angle),
194 gender with meniscus parameters, such as meniscal height and meniscal
195 subluxation were carried out using correlation coefficients (spearman) for each
196 knee. If $r < 0.3$, it was regarded as week correlation, if $r > 0.3$ and < 0.7 , it was
197 regarded as intermediate correlation, and if $r > 0.7$, it was regarded as strong
198 correlation between variables.

199 All analyses were performed using SPSS 11.0 (SPSS Inc., Chicago, IL, USA).
200 The p values of 0.05 or less were considered significant.

201

202 **Results**

203 26 patients were men and 107 were women (ages ranged from 56 to 81,
204 mean age 67.4 ± 6.5 years). The average patient weight was 60.3 ± 9.6 kg
205 (range; 42-93), average body mass index (BMI) was 25.2 ± 3.4 (range; 17.2–
206 32.4), and average height was 154.5 ± 8.5 cm (range; 140–173).

207 **Meniscal morphology**

208 The meniscal morphology for the study sample is summarized in Table 1.
209 The most frequent morphology observed was complete maceration or complete
210 resection (Grade 7) in the anterior horn (42.4%), mid-body (53.8%), and in the
211 posterior horn (52.5%) of the lateral meniscus.

212

213 **Meniscal position**

214 The mean values for subluxation and height of the medial and lateral
215 meniscus for the study patients are summarized in Table 2. The anterior horn of
216 the medial meniscus in cases with macerated lateral meniscus showed
217 significantly more subluxation than that in non-macerated lateral meniscus. The
218 height of midbody of medial meniscus in cases with macerated lateral meniscus
219 was significantly smaller than that in non-macerated lateral meniscus.

220 In the analysis of cases only with non-macerated lateral meniscus, the anterior
221 horn and midbody of lateral meniscus showed significantly more subluxation
222 than that of medial meniscus in same cases Also, the height of the anterior

223 horn and midbody of the lateral meniscus had a significantly smaller size than
224 the medial meniscus in same cases.

225 **FTA angle**

226 The FTA valgus angle was not related with each meniscal morphological grade
227 at anterior and posterior horn. However, in terms of mid-body, FTA angle
228 showed differences between grade 5 and grade 2 ($p=0.030$), between grade 5
229 and grade 4 ($p=0.012$) and between grade 5 and grade 6 ($p=0.018$) (Table 3).

230 Gender was not found to be correlated with the meniscal position, and FTA
231 except subluxation of the midbody of the medial meniscus (Table 4). The BMI
232 was not found to be correlated with the meniscal position and FTA except
233 subluxation of the midbody of the lateral meniscus
234 ($r=0.232, p=0.003$), subluxation of the anterior horn of the medial meniscus
235 ($r=0.223, p=0.005$), and subluxation of the midbody of the medial meniscus
236 ($r=0.257, p=0.001$)

237

238 **Discussion**

239 The results of this study demonstrate that the majority of study patients
240 had a completely macerated or destroyed meniscus with end stage lateral
241 osteoarthritis of the knee; in contrast to our previous study [20] that showed that
242 most cases with advanced medial osteoarthritis had a hypertrophied medial
243 meniscus. These findings suggest that the degenerative changes of the lateral
244 meniscus might progress to a meniscal tear, which can ultimately lead to
245 complete destruction, unlike the medial meniscus. According to the modified
246 WORMS classification, generally the predominant type of injured lateral

247 meniscus in end stage lateral OA was grade 7 (complete maceration/destruction
248 or complete resection) followed by grade 4 (displaced tear or partial resection).
249 In terms of the dimensional changes of the anterior horn of the lateral meniscus,
250 the proportion of grade 4 and grade 7(Grade 3; 4; 5; 7 = 13.9%, 32.9%, 2.5%,
251 42.4%) was higher and grade 3 and grade 5 was lower , as compared to end
252 staged isolated medial osteoarthritis (Grade 3; 4; 5; 7 = 20.4%; 15.6%; 30%;
253 0%). The tear of anterior horn of lateral meniscus were a very common finding
254 (82.9%, 131/158 cases) in lateral compartment osteoarthritis, unlike anterior
255 horn of medial meniscus in end staged isolated medial osteoarthritis [20], which
256 showed incidence of 47.9%(80/167 cases). Regarding the mid-body of the
257 lateral meniscus, the proportion of grade 7 (53.8%) was much higher than
258 (7.2%) for end stage medial osteoarthritis. The overall incidence of midbody
259 tears of lateral meniscus was 94.3%(149/158 cases) in comparison to
260 95.7(160/167 cases) of medial meniscus mid-body of end staged medial
261 osteoarthritis [20].

262 For the posterior horn, the proportion of grade 6 (1.3%) was low and grade 7
263 (53.9%) was much higher, compared to end stage medial osteoarthritis (Grade
264 6, 7 = 83.8%; 0.5%). The overall incidence of post horn damage of lateral
265 meniscus was 94.9%(150/158 cases) in comparison to 98.8%(165/167 cases)
266 of medial meniscus posterior horn of end staged osteoarthritis[20]. These
267 findings indicate that most lateral menisci in persons with end-stage lateral OA
268 are predominantly macerated. However, all parts of the meniscus were not
269 completely macerated; 57.6% (91/158 cases), 46.2% (63/158 cases) and 47.5%
270 (75/158 cases) of each portion of the lateral meniscus was not macerated.

271 Therefore, although the existing dogma appears to be correct in suggesting that
272 in the vast majority of persons with end-stage OA the meniscus is destroyed/
273 macerated, it is important to consider that the entire lateral meniscus was not
274 affected by the same mechanism. In addition, various factors influencing the
275 mechanisms associated with lateral osteoarthritis remain unknown.

276 Limb alignment had a significant association with meniscal morphology of
277 midbody of lateral meniscus, which was high valgus alignment of $17.7 \pm 3.7^\circ$ in
278 grade 5, relative to other grades. These findings are different from those of end
279 stage medial osteoarthritis in previous published our study, which limb
280 alignment was not associated with meniscal morphology[20]. However, as the
281 number of cases of grade 5 was small, further study with larger number of
282 cases might be needed to conclude whether there are definite differences
283 between FTA and meniscal morphology or not. In terms of meniscal position,
284 both non-macerated and macerated lateral menisci were accompanied by the
285 subluxation of the medial meniscus of the same knee. In detail, the anterior
286 horn and midbody of medial meniscus in cases with non-macerated lateral
287 meniscus showed subluxation of 1.30 ± 1.72 mm and 2.44 ± 2.14 mm, respectively.
288 For case of the macerated lateral meniscus, those are 2.77 ± 2.55 mm and 2.26
289 ± 1.98 mm. This finding indicates that the lateral TF osteoarthritis can affect the
290 medial compartment.

291 In terms of meniscal height, lateral meniscus with non-macerated morphology
292 (anterior horn= 5.41 ± 1.98 mm, midbody= 6.03 ± 1.88 mm) showed smaller height
293 than medial meniscus of same cases (anterior horn= 6.35 ± 1.71 mm, midbody
294 7.05 ± 1.83 mm). These finding may be due to the fact that lateral TF

295 osteoarthritis did not have many case of grade 5 and grade 6 than the medial
296 TF osteoarthritis. But this was the case in anterior horn and midbody, not in
297 posterior horn.(Posterior horn of lateral meniscus = 5.66 ± 1.63 , Posterior horn of
298 medial meniscus = 5.89 ± 0.92)

299 The limitations of this study include the following. First, the cohort size
300 (158 knees) was relatively small and different results might have been obtained
301 with a larger study sample. However, based on the low prevalence of lateral
302 osteoarthritis, which is one-tenth that of medial osteoarthritis [26], the results
303 suggest the need for additional research. Second,our study finding can not be
304 generalized to all lateral OA because this study is highly selected sample of
305 severe symptomatic lateral OA scheduled for TKA with no trauma background
306 which is likely not representative for lateral knee OA at large. Third, there is the
307 possibility that the menisci in the subjects might continue to change and
308 become completely destroyed or macerated, which would affect the findings
309 and interpretation of outcomes. Forth, it is unclear whether the hypertrophied
310 lateral menisci in this series (13 cases at the anterior horn, 11 cases at the
311 mid-body, 5 cases at the posterior horn) were truly hypertrophied or
312 alternatively the result of destroyed discoid lateral menisci, a common finding in
313 Korea [27-28]. This would depend on the enrollment of patients with a discoid
314 meniscus in the study, which is unknown and would likely lead to different
315 results. Fifth, as histologic analysis were not done in this series, we cannot
316 definitely conclude that lateral compartment osteoarthritis has various grades of
317 meniscal morphology but we believe that our classification based on previous
318 reports [20] can contribute to understanding different mechanisms of medial

319 and lateral osteoarthritis.

320 Based on the high prevalence of a hypertrophied medial meniscus in patients
321 with end stage medial osteoarthritis, and the high prevalence of a macerated
322 lateral meniscus in patients with end stage lateral osteoarthritis, other factors
323 such as local biomechanical dynamics and the different surrounding structures
324 for each compartment should be considered and investigated to better
325 understand the development and progression of medial and lateral osteoarthritis.
326 Since the macerated meniscus is the final form of lateral OA, the morphological
327 changes of the meniscus in prior grades of end stage lateral OA should be
328 further studied. Based on the results of this study, the final meniscus
329 abnormality of end stage lateral OA appears to be complete maceration.

330

331 **Author contributions**

332 All authors made substantial contributions to all of the following: (1) the
333 conception and design of the study, or acquisition of data, or analysis and
334 interpretation of data, (2) drafting the article or revising it critically for important
335 intellectual content, (3) final approval of the version to be submitted.

336

337 **Conflict of interest**

338 The authors have declared no conflicts of interest.

339

340 **Acknowledgements**

341 The authors would like to thank the staff and participants in the Joint & Arthritis
342 Research of Himchan Hospital.

343

344 **References**

345

- 346 1. Resnick D NG. Degenerative disease. Diagnosis of bone an joint
347 disorders. 2nd edn. Saunders, Philadelphia 1988:1442-58.
- 348 2. Ahlback S. Osteoarthritis of the knee. A radiographic investigation. Acta
349 Radiol Diagn (Stockh) 1968:Suppl 277:7-72.
- 350 3. Thomas RH, Resnick D, Alazraki NP, Daniel D, Greenfield R.
351 Compartmental evaluation of osteoarthritis of the knee. A comparative
352 study of available diagnostic modalities. Radiology 1975;116:585-94.
- 353 4. Sabiston CP, Adams ME, Li DK. Magnetic resonance imaging of
354 osteoarthritis: correlation with gross pathology using an experimental
355 model. J Orthop Res 1987;5:164-72.
- 356 5. Pataki A, Fries R, Ochsner K, Witzemann E. Qualitative radiographic
357 diagnosis of osteo-arthritis of the knee joint in the C57BL mouse. Agents
358 Actions 1987;22:123-30.
- 359 6. Tardif G, Hum D, Pelletier JP, Duval N, Martel-Pelletier J. Regulation of
360 the IGFBP-5 and MMP-13 genes by the microRNAs miR-140 and miR-
361 27a in human osteoarthritic chondrocytes. BMC Musculoskelet Disord
362 2009;10:148.
- 363 7. Ikegawa S. [Progress of research in osteoarthritis. Recent advance in the
364 study of susceptibility genes for osteoarthritis]. Clin Calcium
365 2009;19:1616-20.
- 366 8. Kawaguchi H. [Progress of research in osteoarthritis. Molecular targeting
367 for osteoarthritis treatment]. Clin Calcium 2009;19:1608-14.
- 368 9. Okazaki K, Iwamoto Y. [Progress of research in osteoarthritis. Gene
369 expression and its regulatory mechanisms in the degenerative cartilage
370 in osteoarthritis]. Clin Calcium 2009;19:1578-85.
- 371 10. Wei L, Svensson O, Hjerpe A. Correlation of morphologic and
372 biochemical changes in the natural history of spontaneous osteoarthritis
373 in guinea pigs. Arthritis Rheum 1997;40:2075-83.
- 374 11. Hunter DJ, Zhang YQ, Niu JB, Tu X, Amin S, Clancy M, et al. The
375 association of meniscal pathologic changes with cartilage loss in
376 symptomatic knee osteoarthritis. Arthritis Rheum 2006;54:795-801.
- 377 12. Hunter DJ, Zhang YQ, Tu X, Lavalley M, Niu JB, Amin S, et al. Change
378 in joint space width: hyaline articular cartilage loss or alteration in
379 meniscus? Arthritis Rheum 2006;54:2488-95.

- 380 13. Buck RJ, Wyman BT, Le Graverand MP, Wirth W, Eckstein F. An
381 efficient subset of morphological measures for articular cartilage in the
382 healthy and diseased human knee. *Magn Reson Med* 2010;63:680-90.
- 383 14. Wang SZ, Huang YP, Saarakkala S, Zheng YP. Quantitative assessment
384 of articular cartilage with morphologic, acoustic and mechanical
385 properties obtained using high-frequency ultrasound. *Ultrasound Med
386 Biol* 2010;36:512-27.
- 387 15. Adams JG, McAlindon T, Dimasi M, Carey J, Eustace S. Contribution of
388 meniscal extrusion and cartilage loss to joint space narrowing in
389 osteoarthritis. *Clin Radiol* 1999;54:502-6.
- 390 16. Gale DR, Chaisson CE, Totterman SM, Schwartz RK, Gale ME, Felson
391 D. Meniscal subluxation: association with osteoarthritis and joint space
392 narrowing. *Osteoarthritis Cartilage* 1999;7:526-32.
- 393 17. Erbagci H, Gumusburun E, Bayram M, Karakurum G, Sirikci A. The
394 normal menisci: in vivo MRI measurements. *Surg Radiol Anat*
395 2004;26:28-32.
- 396 18. Ding C, Martel-Pelletier J, Pelletier JP, Abram F, Raynauld JP, Cicuttini F,
397 et al. Meniscal tear as an osteoarthritis risk factor in a largely non-
398 osteoarthritic cohort: a cross-sectional study. *J Rheumatol* 2007;34:776-
399 84.
- 400 19. Englund M, Guermazi A, Gale D, Hunter DJ, Aliabadi P, Clancy M, et al.
401 Incidental meniscal findings on knee MRI in middle-aged and elderly
402 persons. *N Engl J Med* 2008;359:1108-15.
- 403 20. Jung KA, Lee SC, Hwang SH, Yang KH, Kim DH, Sohn JH, et al. High
404 frequency of meniscal hypertrophy in persons with advanced varus knee
405 osteoarthritis. *Rheumatol Int* 2010;30:1325-33.
- 406 21. Wirth W, Frobell RB, Souza RB, Li X, Wyman BT, Le Graverand MP, et
407 al. A three-dimensional quantitative method to measure meniscus shape,
408 position, and signal intensity using MR images: a pilot study and
409 preliminary results in knee osteoarthritis. *Magn Reson Med*
410 2010;63:1162-71.
- 411 22. Felson DT, Nevitt MC, Zhang Y, Aliabadi P, Baumer B, Gale D, et al.
412 High prevalence of lateral knee osteoarthritis in Beijing Chinese
413 compared with Framingham Caucasian subjects. *Arthritis Rheum*
414 2002;46:1217-22.
- 415 23. Kellgren JH LJ. *Atlas of standard radiographs*. Blackwell, Oxford 1963.

416 24. Altman RD, Hochberg M, Murphy WA, Jr., Wolfe F, Lequesne M. Atlas of
417 individual radiographic features in osteoarthritis. *Osteoarthritis Cartilage*
418 1995;3 Suppl A:3-70.

419 25. Peterfy CG, Guermazi A, Zaim S, Tirman PF, Miaux Y, White D, et al.
420 Whole-Organ Magnetic Resonance Imaging Score (WORMS) of the
421 knee in osteoarthritis. *Osteoarthritis Cartilage* 2004;12:177-90.

422 26. Scott RD. Lateral unicompartmental replacement: a road less traveled.
423 *Orthopedics* 2005;28:983-4.

424 27. Kim SJ, Lee YT, Kim DW. Intraarticular anatomic variants associated
425 with discoid meniscus in Koreans. *Clin Orthop Relat Res* 1998:202-7.

426 28. Seong SC, Park MJ. Analysis of the discoid meniscus in Koreans.
427 *Orthopedics* 1992;15:61-5.

428

429 .

430

431

432

433

434

435

436

437

438 **Legends**

439 **Fig. 1 A, B.** The modification of the WORMS assessment method²⁴.

440 Schematic drawing (a) and MRI findings (b) are representative of the global
441 meniscus scoring system.

442 0= intact,

443 1= minor radial tear or parrot-beak tear

444 2= nondisplaced tear

445 3= displaced but no tear

446 4= displaced tear or partial resection

447 5= hypertrophied displaced


448 6= hypertrophied displaced tear

449 7=complete maceration/destruction or complete resection

450 “Hypertrophy” > 2 mm larger than MM


451

452 **A)**



453

454 **B)**



455

456


457

458

459

460 **Fig. 2 A, B.** Meniscal position was assessed by measuring meniscal
461 subluxation and height for each knee. To determine the meniscal height, the
462 anterior and posterior horns of the menisci were measured in the sagittal plane,
463 which allowed for the best visualization of the greatest meniscal size. The mid-
464 body height was measured in the coronal plane, where the medial and lateral
465 tibial spine volume was maximal. (a) The meniscal height was measured at the
466 most peripheral edge of each meniscus, regardless of whether the meniscus
467 was “in-place”, subluxed or extruded. To determine the meniscal subluxation,
468 the anterior subluxation of the anterior horn of the medial and lateral meniscus
469 was assessed in the area where the subluxation was most prominent, based on
470 multiple sagittal slices.(b) Lateral subluxation of the mid-body of the lateral and
471 medial subluxation of the medial meniscus was measured where the volume of
472 the medial and lateral tibial spine was greatest.* Posterior subluxation of the
473 posterior horn was not measured, because this could not be performed
474 accurately in the sagittal plane. For the menisci that were completely macerated
475 or destroyed, meniscal subluxation and meniscal height could not be measured.

476
477
478
479
480
481
482
483
484
485
486



487
488
489

Table 1. Meniscal morphology at each portion of the lateral meniscus assessed by a modified WORMS method

	Grade 0(%)	Grade 1(%)	Grade 2(%)	Grade 3(%)	Grade 4(%)	Grade 5(%)	Grade 6(%)	Grade 7(%)	Total (%)
Ant. Horn of LM	1(0.6)	0	3(1.9)	22(13.9)	52(32.9)	4(2.5)	9(5.7)	67(42.4)*	158(100)
Mid-body of LM	1(0.6)	1(0.6)	6(3.8)	4(2.5)	50(31.6)	4(2.5)	7(4.4)	85(53.8)*	158(100)
Post. Horn of LM	1(0.6)	0	1(0.6)	5(3.2)	63(39.9)	2(1.3)	3(1.9)	83(52.5)*	158(100)

Grade 0 = intact, 1 = minor radial tear or parrot-beak tear, 2 = nondisplaced tear, 3 = displaced but no tear, 4 = displaced tear or partial resection, 5 = hypertrophied displaced 6 = hypertrophied displaced tear, 7 = complete maceration/destruction or complete resection

*Predominant type in each portion

Table 2. Meniscal position and Meniscal Height for each region of the meniscus, as determined by MR imaging.
MM= Medial meniscus, LM= Lateral meniscus Mean BMI : 25.8 +_ 3.36(Range:17.92 to 32.39)

		No. of knees (n, total n = 158)	Subluxation of LM	Suluxation of MM	Height of LM	Height of MM
Macerated LM (Grade 7)	Ant. horn	67	Not measured	2.77±2.55 mm (0 to 12.31) [§]	Not measured	5.38 ± 1.10 mm (3.44 to 8.50)
	Midbody	85	Not measured	2.26±1.98 mm (0.0 to 7.91)	Not measured	5.92 ± 1.60 mm (2.93 to 9.67)
	Post. Horn	83	Not measured	Not measured	Not measured	5.56 ± 1.40 mm (2.5 to 8.9)
Non-macerated LM (remainder)	Ant. horn	91	6.45 ± 3.11 mm* (0 to 12.01)	1.30±1.72mm (0 to 6.74)	5.41 ± 1.98 mm (2.03 to 11.15)	6.35 ± 1.71 mm** (3.01 to 7.91)
	Midbody	73	5.27 ±3.37 mm* (0 to 14.94)	2.44±2.14mm (0 to 7.32)	6.03 ± 1.88 mm (0.25 to 11.70)	7.05 ± 1.83 mm** ^{§§} (3.14 to 12.89)
	Post. Horn	75	Not measured	Not measured	5.66 ± 1.63 mm (3.01 to 9.08)	5.89 ± 0.92 mm (4.1 to 7.62)

MM medial meniscus, LM lateral meniscus, n total number of knee,

* Statistically significant difference between LM subluxation and MM subluxation for the cases with non-macerated LM, using independent T-test at the anterior horn and midbody (p=0.001).

** Statistically significant difference between LM height and MM height for the cases with non-macerated LM, using independent T-test at the anterior horn and midbody (p=0.001).

[§] Statistically significant difference in MM subluxation between cases with macerated LM and non-macerated LM, using independent T-test at the anterior horn (p=0.0001).

^{§§} Statistically significant difference in MM height between cases with macerated LM and non-macerated LM, using independent T-test at the anterior horn and midbody (p=0.003, p=0.002).

Table 3. Association between FTA and each meniscal morphology

Meniscus morphology	Anterior horn of lateral meniscus *		Midbody of lateral meniscus **		Posterior horn of lateral meniscus ***	
	No.of knee	FTA mean±SD	No.of knee	FTA mean±SD	No.of knee	FTA mean±SD
0	1	8.15	1	0.5	1	0.5
1	0	-	1	2.8	0	-
2	3	11.8±6.4	6	7.2±2.6	1	2.8
3	22	8.2±4.5	4	13.0±4.5	5	13.1±5.8
4	52	8.0±4.6	49	8.4±4.5	62	8.8±5.7
5	4	13.0±9.8	4	17.7±3.7	2	16.3±1.3
6	9	9.1±7.0	7	7.0±3.9	3	6.3±0.9
7	66	10.6±5.8	86	10.0±5.9	84	9.7±5.3
Total	158	9.4±5.5	158	9.4±5.5	158	9.4±5.5

* anterior horn : P-value (oneway ANOVA test): 0.067

** midbody: P-value (oneway ANOVA test): 0.007, grade 5 Vs grade 2 (P=0.030), grade 5 Vs grade 4 (p=0.012). grade 5 Vs grade 6 (p=0.018) in Post-hoc test(Tukey test)

*** posterior horn : P-value (oneway ANOVA test): 0.113

Table 4 Association between sex and meniscal subluxation ,FTA

	Sex	No. of Knees	Mean \pm SD	P value
FTA	F	126	9.15 \pm 5.36	0.267
	M	32	10.35 \pm 5.89	
Subluxation of LM (ant horn)	F	126	3.56 \pm 3.93	0.335
	M	32	4.32 \pm 4.12	
Subluxation of LM (midbody)	F	126	2.44 \pm 3.61	0.180
	M	32	3.40 \pm 3.54	
Subluxation of MM (ant horn)	F	126	1.85 \pm 2.19	0.460
	M	32	2.18 \pm 2.37	
Subluxation of MM (midbody)	F	126	2.51 \pm 2.03	0.042
	M	32	1.68 \pm 2.01	